

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/00	A2	(11) International Publication Number: WO 00/21986 (43) International Publication Date: 20 April 2000 (20.04.00)
(21) International Application Number: PCT/US99/23315 (22) International Filing Date: 6 October 1999 (06.10.99) (30) Priority Data: 09/169,289 9 October 1998 (09.10.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/169,289 (CIP) Filed on 9 October 1998 (09.10.98) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): WALKER, Michael, G. [CA/US]; Unit 80, 1050 Borregas Avenue, Sunnyvale, CA 94089 (US). VOLKMUTH, Wayne [US/US]; 783 Roble Avenue, #1, Menlo Park, CA 94025 (US). KLINGLER, Tod, M. [US/US]; 28 Dover Court, San Carlos, CA 94070 (US). (74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: MATRIX-REMODELING GENES (57) Abstract The invention provides novel matrix-remodeling genes and polypeptides encoded by those genes. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing diseases associated with matrix remodeling.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

MATRIX-REMODELING GENES

TECHNICAL FIELD

5 The invention relates to novel matrix-remodeling genes identified by their coexpression with known matrix-remodeling genes. The invention also relates to the use of these biomolecules in diagnosis, prognosis, prevention, treatment, and evaluation of therapies for diseases, particularly diseases associated with matrix-remodeling such as cancer, cardiomyopathy, arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and
10 ulceration.

BACKGROUND OF THE INVENTION

Matrix remodeling is associated with the construction, destruction, and reorganization of extracellular matrix components and is essential in normal cellular functions and also in many disease processes. These disease processes include metastatic cancer, cardiomyopathy, arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and ulceration (Alexander
15 and Werb (1991) In: Cell Biology of Extracellular Matrix, Plenum Press, New York NY, pp. 255-302; Schuppan et al. (1993) In: Extracellular Matrix, Marcel Dekker, New York NY, pp. 201-254; Zvibel and Kraft (1993) In: Extracellular Matrix, Marcel Dekker, New York NY, pp. 559-580; Shanahan et al. (1994) J Clin Invest 93:2393-402; Kielty and Shuttleworth (1995) Int
20 J Biochem Cell Biol 27:747-60; Bitar and Labbad (1996) J Surg Res 61:113-9; Dourado et al. (1996) Osteoarthritis Cartilage 4:187-96; Grant et al. (1996) Regul. Pept. 67:137-44; Gunja-Smith et al. (1996) Am J Pathol 148:1639-48; Alcolado et al. (1997) Clin. Sci 92:103-12; Cs-Szabo et al. (1997) Arthritis Rheum 40:1037-45; Hayward and Brock (1997) Hum Mutat 10:415-23; Ledda et al. (1997) J Invest Dermatol 108:210-4; Hayashido et al. (1998) Int J
25 Cancer 75:654-8; Ito et al. (1998) Kidney Int 53:853-61; Nelson et al. (1998) Cancer Res 58:232-6).

Many genes that participate in and regulate matrix remodeling are known, but many remain to be identified. Identification of currently unknown genes will provide new diagnostic and therapeutic targets. In addition, these genes will provide new opportunities for
30 therapeutic tissue engineering--the use of drugs or biologicals to direct the creation of new tissues such as skin, pancreas, or liver that can replace tissues lost to disease or trauma.

The present invention provides new compositions that are useful for diagnosis, prognosis, treatment, prevention, and evaluation of therapies for diseases associated with matrix remodeling. We have implemented a method for analyzing gene expression patterns and have identified 20 novel matrix-remodeling genes by their coexpression with known
5 matrix-remodeling genes.

SUMMARY OF THE INVENTION

In one aspect, the invention provides for a substantially purified polynucleotide comprising a gene that is coexpressed with one or more known matrix-remodeling genes in a plurality of biological samples. Preferably, each known matrix-remodeling gene is selected
10 from the group consisting of osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, III, and IV, connective tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptor (fibr-r), fibulin 1, heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor 1 (IGF 1), insulin-like growth factor binding protein (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix
15 metalloproteases (MMP), and tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3). Preferred embodiments are (a) a polynucleotide sequence selected from the group consisting of SEQ ID NOs:1-20; (b) a polynucleotide sequence which encodes a polypeptide sequence of SEQ ID NOs:21, 22, and 23; (c) a polynucleotide sequence having at least 70% identity to the polynucleotide sequence of (a) or (b); (d) a polynucleotide
20 sequence comprising at least 18 sequential nucleotides of the polynucleotide sequence of (a), (b), or (c); (e) a polynucleotide which hybridizes under stringent conditions to the polynucleotide of (a), (b), (c), or (d); or (f) a polynucleotide sequence which is complementary to the polynucleotide sequence of (a), (b), (c), (d) or (e). Furthermore, the invention provides an expression vector comprising any of the above described
25 polynucleotides and host cells comprising the expression vector. Still further, the invention provides a method for treating or preventing a disease or condition associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a sample comprising administering to a subject in need the above-described polynucleotides in an amount effective for treating or preventing said disease.

30 In a second aspect, the invention provides a substantially purified polypeptide comprising the gene product of a gene that is coexpressed with one or more known matrix-

remodeling genes in a plurality of biological samples. The known matrix-remodeling gene may be selected from the group consisting of osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, II, and IV, connective tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptors (fibr-r), fibulin 1, heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor 1 (IGF 1), insulin-like growth factor binding protein (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix metalloproteases (MMP), and tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3). Preferred embodiments are polypeptides comprising (a) the polypeptide sequence of SEQ ID NO:21, 22, or 23; (b) a polypeptide sequence having at least 85% identity to the polypeptide sequence of (a); and (c) a polypeptide sequence comprising at least 6 sequential amino acids of the polypeptide sequence of (a) or (b). Additionally, the invention provides antibodies that bind specifically to any of the above described polypeptides and a method for treating or preventing a disease or condition associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a sample comprising administering to a subject in need such an antibody in an amount effective for treating or preventing said disease.

In another aspect, the invention provides a pharmaceutical composition comprising the polynucleotide of claim 2 or the polypeptide of claim 3 in conjunction with a suitable pharmaceutical carrier or a method for treating or preventing a disease or condition associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a sample comprising administering to a subject in need such compositioning in an amount effective for treating or preventing said disease.

In yet a further aspect, the invention provides a method for diagnosing a disease or condition associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a sample, wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, II, and IV, connective tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptor (fibr-r), fibulin 1, heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor 1 (IGF 1), insulin-like growth factor binding protein (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix metalloproteases (MMP), and tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP

- 1, 2, and 3). The method comprises the steps of (a) providing the sample comprising one of more of said coexpressed genes; (b) hybridizing the polynucleotide of the coexpressed genes under conditions effective to form one or more hybridization complexes; and (c) detecting the hybridization complexes, wherein the altered level of one or more of the hybridization
- 5 complexes in a diseased sample compared with the level of hybridization complexes in a non-diseased sample correlates with the presence of the disease or condition in the sample.

BRIEF DESCRIPTION OF THE SEQUENCE LISTING

The Sequence Listing provides exemplary matrix-remodeling-associated sequences including polynucleotide sequences, SEQ ID NOs:1-20, and polypeptide sequences, SEQ ID

10 NOs:21-23. Each sequence is identified by a sequence identification number (SEQ ID NO) and by the Incyte Clone number from which the sequence was first identified.

DESCRIPTION OF THE INVENTION

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include the plural reference unless the context clearly dictates otherwise.

15 Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

DEFINITIONS

"NSEQ" refers generally to a polynucleotide sequence of the present invention,

20 including SEQ ID NOs:1-20. "PSEQ" refers generally to a polypeptide sequence of the present invention, including SEQ ID NOs:21-23.

A "variant" refers to either a polynucleotide or a polypeptide whose sequence diverges from SEQ ID NOs:1-20 or SEQ ID NOs:21-23, respectively. Polynucleotide sequence divergence may result from mutational changes such as deletions, additions, and substitutions

25 of one or more nucleotides; it may also occur because of differences in codon usage. Each of these types of changes may occur alone, or in combination, one or more times in a given sequence. Polypeptide variants include sequences that possess at least one structural or functional characteristic of SEQ ID NOs:21-23.

A "fragment" can refer to a nucleic acid sequence that is preferably at least 20 nucleic

30 acids in length, more preferably 40 nucleic acids, and most preferably 60 nucleic acids in length, and encompasses, for example, fragments consisting of nucleic acids 1-50 or 200-500

of SEQ ID NOs:1-20. A "fragment" can also refer to polypeptide sequences which are preferably at least 5 to about 15 amino acids in length, most preferably at least 10 amino acids long, and which retain some biological activity or immunological activity of, for example, a sequence selected from SEQ ID NOs:21-23.

- 5 "Gene" or "gene sequence" refers to the partial or complete coding sequence of a transcript. The term also refers to sequences corresponding to 5' or 3' untranslated regions or 5' or 3' untranslated regions including partial or complete coding sequences of a gene. Typically, the novel gene sequences may or may not be homologous to annotated sequences found in public or private databases. The gene may be in a sense or antisense
10 (complementary) orientation.

"Known matrix-remodeling gene" refers to a gene sequence which has been previously identified as useful in the diagnosis, treatment, prognosis, or prevention of diseases associated with matrix remodeling. Typically, this means that the known matrix-remodeling gene is expressed at higher levels in tissue abundant in known matrix-remodeling transcripts when
15 compared with other tissue.

"Matrix-remodeling gene" refers to a gene sequence whose expression pattern is similar to that of the known matrix-remodeling genes and which are useful in the diagnosis, treatment, prognosis, or prevention of diseases associated with matrix remodeling. The gene sequences can also be used in the evaluation of therapies for cancer.

- 20 "Substantially purified" refers to a nucleic acid or an amino acid sequence that is removed from its natural environment and is isolated or separated, and is at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which it is naturally present.

THE INVENTION

- 25 The present invention encompasses a method for identifying biomolecules that are associated with a specific disease, regulatory pathway, subcellular compartment, cell type, tissue type, or species. In particular, the method identifies gene sequences useful in diagnosis, prognosis, treatment, prevention, and evaluation of therapies for diseases associated with matrix-remodeling, particularly, cancer, cardiomyopathy, arthritis, angiogenesis, diabetic
30 necrosis, atherosclerosis, fibrosis, and ulceration.

The method provides first identifying polynucleotides that are expressed in a plurality

of cDNA libraries. The identified polynucleotides include genes of known function, genes known to be specifically expressed in a specific disease process, subcellular compartment, cell type, tissue type, or species. Additionally, the polynucleotides include genes of unknown function. The expression patterns of the known genes are then compared with those of the
5 genes of unknown function to determine whether a specified coexpression probability threshold is met. Through this comparison, a subset of the polynucleotides for unknown function genes having a high coexpression probability with the known genes can be identified. The high coexpression probability correlates with a particular coexpression probability threshold which is less than 0.001, and more preferably less than 0.00001.

10 The polynucleotides originate from cDNA libraries derived from a variety of sources including, but not limited to, eukaryotes such as human, mouse, rat, dog, monkey, plant, and yeast and prokaryotes such as bacteria and viruses. These polynucleotides can also be selected from a variety of sequence types including, but not limited to, expressed sequence tags (ESTs), assembled polynucleotide sequences, full length gene coding regions, introns,
15 regulatory sequences, 5' untranslated regions, and 3' untranslated regions. To have statistically significant analytical results, the polynucleotides need to be expressed in at least three cDNA libraries.

The cDNA libraries used in the coexpression analysis of the present invention can be obtained from blood vessels, heart, blood cells, cultured cells, connective tissue, epithelium,
20 islets of Langerhans, neurons, phagocytes, biliary tract, esophagus, gastrointestinal system, liver, pancreas, fetus, placenta, chromaffin system, endocrine glands, ovary, uterus, penis, prostate, seminal vesicles, testis, bone marrow, immune system, cartilage, muscles, skeleton, central nervous system, ganglia, neuroglia, neurosecretory system, peripheral nervous system, bronchus, larynx, lung, nose, pleurus, ear, eye, mouth, pharynx, exocrine glands, bladder,
25 kidney, ureter, and the like. The number of cDNA libraries selected can range from as few as 20 to greater than 10,000. Preferably, the number of the cDNA libraries is greater than 500.

In a preferred embodiment, gene sequences are assembled to reflect related sequences, such as assembled sequence fragments derived from a single transcript. Assembly of the polynucleotide sequences can be performed using sequences of various types including, but
30 not limited to, ESTs, extensions, or shotgun sequences. In a most preferred embodiment, the polynucleotide sequences are derived from human sequences that have been assembled using

the algorithm disclosed in "Database and System for Storing, Comparing and Displaying Related Biomolecular Sequence Information", Lincoln et al., Serial No:60/079,469, filed March 26, 1998, incorporated herein by reference.

Experimentally, differential expression of the polynucleotides can be evaluated by
5 methods including, but not limited to, differential display by spatial immobilization or by gel electrophoresis, genome mismatch scanning, representational difference analysis, and transcript imaging. Additionally, differential expression can be assessed by microarray technology. These methods may be used alone or in combination.

Known matrix-remodeling genes can be selected based on the use of the genes as
10 diagnostic or prognostic markers or as therapeutic targets for diseases associated with matrix remodeling, such as cancer, cardiomyopathy, arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and ulceration. Preferably, the known matrix-remodeling genes include osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, III, and IV, connective tissue growth factor (CTGF), fibrillin, fibronectins,
15 fibronectin receptor (fibr-r), fibulin 1, heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor 1 (IGF 1), insulin-like growth factor binding protein (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix metalloproteinases (MMP), tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3), and the like.

20 The procedure for identifying novel genes that exhibit a statistically significant coexpression pattern with known matrix-remodeling genes is as follows. First, the presence or absence of a gene sequence in a cDNA library is defined: a gene is present in a cDNA library when at least one cDNA fragment corresponding to that gene is detected in a cDNA sample taken from the library, and a gene is absent from a library when no corresponding cDNA
25 fragment is detected in the sample.

Second, the significance of gene coexpression is evaluated using a probability method to measure a due-to-chance probability of the coexpression. The probability method can be the Fisher exact test, the chi-squared test, or the kappa test. These tests and examples of their applications are well known in the art and can be found in standard statistics texts (Agresti, A
30 (1990) Categorical Data Analysis, John Wiley & Sons, New York NY; Rice, JA (1988) Mathematical Statistics and Data Analysis, Duxbury Press, Pacific Grove CA). A Bonferroni

correction (Rice, supra, page 384) can also be applied in combination with one of the probability methods for correcting statistical results of one gene versus multiple other genes. In a preferred embodiment, the due-to-chance probability is measured by a Fisher exact test, and the threshold of the due-to-chance probability is set to less than 0.001, more preferably less than 0.00001.

To determine whether two genes, A and B, have similar coexpression patterns, occurrence data vectors can be generated as illustrated in Table 1, wherein a gene's presence is indicated by a one and its absence by a zero. A zero indicates that the gene did not occur in the library, and a one indicates that it occurred at least once.

Table 1. Occurrence data for genes A and B

	Library 1	Library 2	Library 3	...	Library N
gene A	1	1	0	...	0
gene B	1	0	1	...	0

For a given pair of genes, the occurrence data in Table 1 can be summarized in a 2 x 2 contingency table.

Table 2. Contingency table for co-occurrences of genes A and B

	Gene A present	Gene A absent	Total
Gene B present	8	2	10
Gene B absent	2	18	20
Total	10	20	30

Table 2 presents co-occurrence data for gene A and gene B in a total of 30 libraries. Both gene A and gene B occur 10 times in the libraries. Table 2 summarizes and presents 1) the number of times gene A and B are both present in a library, 2) the number of times gene A and B are both absent in a library, 3) the number of times gene A is present while gene B is absent, and 4) the number of times gene B is present while gene A is absent. The upper left entry is the number of times the two genes co-occur in a library, and the middle right entry is the number of times neither gene occurs in a library. The off diagonal entries are the number of times one gene occurs while the other does not. Both A and B are present eight times and absent 18 times, gene A is present while gene B is absent two times, and gene B is present while gene A

is absent two times. The probability ("p-value") that the above association occurs due to chance as calculated using a Fisher exact test is 0.0003. Associations are generally considered significant if a p-value is less than 0.01 (Agresti, supra; Rice, supra).

This method of estimating the probability for coexpression of two genes makes several
5 assumptions. The method assumes that the libraries are independent and are identically
sampled. However, in practical situations, the selected cDNA libraries are not entirely
independent because more than one library may be obtained from a single patient or tissue,
and they are not entirely identically sampled because different numbers of cDNAs may be
sequenced from each library (typically ranging from 5,000 to 10,000 cDNAs per library). In
10 addition, because a Fisher exact coexpression probability is calculated for each gene versus
41,419 other genes, a Bonferroni correction for multiple statistical tests is necessary.

Using the method of the present invention, we have identified 20 novel genes that
exhibit strong association, or coexpression, with known genes that are matrix-remodeling-
specific. These known matrix-remodeling genes include osteonectin (BM-40),
15 chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, III, and IV, connective
tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptor (fibr-r), fibulin 1,
heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like
growth factor 1 (IGF 1), insulin-like growth factor binding protein (IGFBP), laminin, lumican,
matrix Gla protein (MGP), matrix metalloproteases (MMP), and tissue inhibitors of matrix
20 metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3). The results presented in Tables 5 and 6
show that the expression of the 20 novel genes have direct or indirect association with the
expression of known matrix-remodeling genes. Therefore, the novel genes can potentially be
used in diagnosis, treatment, prognosis, or prevention of diseases associated with matrix
remodeling, or in the evaluation of therapies for diseases associated with matrix remodeling.
25 Further, the gene products of the 20 novel genes are potential therapeutic proteins and targets
of therapeutics against diseases associated with matrix remodeling.

Therefore, in one embodiment, the present invention encompasses a polynucleotide
sequence comprising the sequence of SEQ ID NOs:1-20. These 20 polynucleotides are shown
by the method of the present invention to have strong coexpression association with known
30 matrix-remodeling genes and with each other. The invention also encompasses a variant of
the polynucleotide sequence, its complement, or 18 consecutive nucleotides of a sequence

provided in the above described sequences. Variant polynucleotide sequences typically have at least about 70%, more preferably at least about 85%, and most preferably at least about 95% polynucleotide sequence identity to NSEQ.

One preferred method for identifying variants entails using NSEQ and/or PSEQ sequences to search against the GenBank primate (pri), rodent (rod), and mammalian (mam), vertebrate (vrtp), and eukaryote (eukp) databases, SwissProt, BLOCKS (Bairoch et al. (1997) *Nucleic Acids Res* 25:217-221), PFAM, and other databases that contain previously identified and annotated motifs, sequences, and gene functions. Methods that search for primary sequence patterns with secondary structure gap penalties (Smith et al. (1992) *Protein Engineering* 5:35-51) as well as algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul (1993) *J Mol Evol* 36:290-300; and Altschul et al. (1990) *J Mol Biol* 215:403-410), BLOCKS (Henikoff and Henikoff (1991) *Nucleic Acids Res* 19:6565-6572), Hidden Markov Models (HMM; Eddy (1996) *Cur Opin Str Biol* 6:361-365; Sonnhammer et al. (1997) *Proteins* 28:405-420), and the like, can be used to manipulate and analyze nucleotide and amino acid sequences. These databases, algorithms and other methods are well known in the art and are described in Ausubel et al. (1997; Short Protocols in Molecular Biology, John Wiley & Sons, New York NY) and in Meyers (1995; Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853).

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to SEQ ID NOS:1-20, and fragments thereof under stringent conditions. Stringent conditions can be defined by salt concentration, temperature, and other chemicals and conditions well known in the art. In particular, stringency can be increased by reducing the concentration of salt, or raising the hybridization temperature. Varying additional parameters, such as hybridization time, the concentration of detergent or solvent, and the inclusion or exclusion of carrier DNA, are well known to those skilled in the art. Additional variations on these conditions will be readily apparent to those skilled in the art (Wahl and Berger (1987) *Methods Enzymol* 152:399-407; Kimmel (1987) *Methods Enzymol* 152:507-511; Ausubel *supra*; and Sambrook et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY).

NSEQ or the polynucleotide sequences encoding PSEQ can be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to

detect upstream sequences, such as promoters and regulatory elements. (See, e.g., Dieffenbach and Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview NY; Sarkar (1993) PCR Methods Applic 2:318-322; Triglia et al. (1988) Nucleic Acids Res 16:8186; Lagerstrom et al. (1991) PCR Methods Applic 1:111-119; and
5 Parker et al. (1991) Nucleic Acids Res 19:3055-306). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto, CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially
10 available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 18 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

In another aspect of the invention, NSEQ or the polynucleotide sequences encoding PSEQ can be cloned in recombinant DNA molecules that direct expression of PSEQ or the
15 polypeptides encoded by NSEQ, or structural or functional fragments thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express the polypeptides of PSEQ or the polypeptides encoded by NSEQ. The nucleotide sequences of the present invention can be engineered using methods
20 generally known in the art in order to alter the nucleotide sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce
25 mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In order to express a biologically active polypeptide encoded by NSEQ, NSEQ or the polynucleotide sequences encoding PSEQ, or derivatives thereof, may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for
30 transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible

promoters, and 5' and 3' untranslated regions in the vector and in NSEQ or polynucleotide sequences encoding PSEQ. Methods which are well known to those skilled in the art may be used to construct expression vectors containing NSEQ or polynucleotide sequences encoding PSEQ and appropriate transcriptional and translational control elements. These methods
5 include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook (supra) and Ausubel (supra)).

A variety of expression vector/host cell systems may be utilized to contain and express NSEQ or polynucleotide sequences encoding PSEQ. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or
10 cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (baculovirus); plant cell systems transformed with viral expression vectors, cauliflower mosaic virus (CaMV) or tobacco mosaic virus (TMV), or with bacterial expression vectors (Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed. For long term production of
15 recombinant proteins in mammalian systems, stable expression of a polypeptide encoded by NSEQ in cell lines is preferred. For example, NSEQ or sequences encoding PSEQ can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector.

20 In general, host cells that contain NSEQ and that express PSEQ may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences. Immunological
25 methods for detecting and measuring the expression of PSEQ using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS).

Host cells transformed with NSEQ or polynucleotide sequences encoding PSEQ may
30 be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly

depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides of NSEQ or polynucleotides encoding PSEQ may be designed to contain signal sequences which direct secretion of PSEQ or polypeptides encoded by NSEQ through a prokaryotic or eukaryotic cell membrane.

5 In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to specify protein targeting, folding,
10 and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant NSEQ or
15 nucleic acid sequences encoding PSEQ are ligated to a heterologous sequence resulting in translation of a fusion protein containing heterologous protein moieties in any of the aforementioned host systems. Such heterologous protein moieties facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin
20 (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, hemagglutinin (HA) and monoclonal antibody epitopes..

In another embodiment, NSEQ or sequences encoding PSEQ are synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers et al. (1980) Nucleic Acids Symp Ser (7) 215-223; Horn et al. (1980) Nucleic Acids Symp Ser (7) 225-232;
25 and Ausubel, supra). Alternatively, PSEQ or a polypeptide sequence encoded by NSEQ itself, or a fragment thereof, may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge et al. (1995) Science 269:202-204). Automated synthesis may be achieved using the ABI 431A Peptide synthesizer (PE Biosystems, Foster City CA). Additionally, PSEQ or the amino acid
30 sequence encoded by NSEQ, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a polypeptide

variant.

In another embodiment, the invention provides a substantially purified polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23 or fragments thereof.

5 **DIAGNOSTICS and THERAPEUTICS**

The sequences of these genes can be used in diagnosis, prognosis, treatment, prevention, and evaluation of therapies for diseases associated with matrix-remodeling, particularly cancer, cardiomyopathy, arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and ulceration. Further, the amino acid sequences encoded by the novel genes are
10 potential therapeutic proteins and targets of anti-cancer therapeutics or for the treatment of other diseases associated with matrix remodeling.

In one preferred embodiment, the polynucleotide sequences of NSEQ or the polynucleotides encoding PSEQ are used for diagnostic purposes to investigate the altered expression of PSEQ, and to monitor regulation of the levels of mRNA or the polypeptides
15 encoded by NSEQ during therapeutic intervention. The polynucleotides may be at least 18 nucleotides long, and may be complementary RNA or DNA molecules, branched nucleic acids, or peptide nucleic acids (PNAs). Alternatively, the polynucleotides are used to detect and quantitate gene expression in samples in which expression of PSEQ or the polypeptides encoded by NSEQ are correlated with disease. Additionally, NSEQ or the polynucleotides
20 encoding PSEQ can be used to detect genetic polymorphisms associated with a disease. These polymorphisms may be detected at the transcript cDNA or genomic level.

The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low), will determine
25 whether the probe identifies only naturally occurring sequences encoding PSEQ, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably have at least 70% sequence identity to any of the NSEQ or PSEQ-encoding sequences.

Means for producing specific hybridization probes for DNAs encoding PSEQ include
30 the cloning of NSEQ or polynucleotide sequences encoding PSEQ into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available,

and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin
5 coupling systems, by fluorescent labels and the like. The polynucleotide sequences encoding PSEQ may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; and in microarrays utilizing fluids or tissues from patients to detect altered NSEQ expression. Such qualitative or quantitative methods are well known in the art.

10 NSEQ or the nucleotide sequences encoding PSEQ can be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value, typically, derived from a non-diseased sample. If the amount of signal in the patient sample is significantly
15 altered in comparison to the standard value then the presence of altered levels of nucleotide sequences of NSEQ and those encoding PSEQ in the sample indicates the presence of the associated disease. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

20 Once the presence of a disease is established and a treatment protocol is initiated, hybridization or amplification assays can be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in a healthy subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

25 The polynucleotides may be used for the diagnosis of a variety of diseases associated with matrix-remodeling including cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary
30 glands, skin, spleen, testis, thymus, thyroid, and uterus, cardiomyopathy, arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and ulceration.

Alternatively, the polynucleotides may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify splice variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disease, to diagnose a disease, and to develop and monitor the activities of therapeutic agents.

In yet another alternative, polynucleotides may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Fluorescent *in situ* hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich et al. (1995) in Meyers, *supra*, pp. 965-968.)

In another embodiment, antibodies which specifically bind PSEQ may be used for the diagnosis of diseases characterized by the over-or-underexpression of PSEQ or polypeptides encoded by NSEQ. A variety of protocols for measuring PSEQ or the polypeptides encoded by NSEQ, including ELISAs, RIAs, and FACS, are well known in the art and provide a basis for diagnosing altered or abnormal levels of the expression of PSEQ or the polypeptides encoded by NSEQ. Standard values for PSEQ expression are established by combining body fluids or cell extracts taken from healthy subjects, preferably human, with antibody to PSEQ or a polypeptide encoded by NSEQ under conditions suitable for complex formation. The amount of complex formation may be quantitated by various methods, preferably by photometric means. Quantities of PSEQ or the polypeptides encoded by NSEQ expressed in disease samples from, for example, biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing or monitoring disease. Alternatively, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PSEQ or the polypeptides encoded by NSEQ specifically compete with a test compound for binding the polypeptides. Antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PSEQ or the polypeptides encoded by NSEQ.

In another aspect, the polynucleotides and polypeptides of the present invention can be employed for treatment or the monitoring of therapeutic treatments for cancer. The polynucleotides of NSEQ or those encoding PSEQ, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotides of NSEQ or those encoding PSEQ may be used in situations in which it would be desirable to

block the transcription or translation of the mRNA.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding PSEQ. (See, e.g., Sambrook, supra; and Ausubel, supra.)

Genes having polynucleotide sequences of NSEQ or those encoding PSEQ can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding PSEQ. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee et al. (1994) In: Huber and Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. Alternatively, nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases may be included.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g.,

Goldman et al. (1997) Nature Biotechnology 15:462-466.)

Further, an antagonist or antibody of a polypeptide of PSEQ or encoded by NSEQ may be administered to a subject to treat or prevent a cancer associated with increased expression or activity of PSEQ. An antibody which specifically binds the polypeptide may be used
5 directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express the the polypeptide.

Antibodies to PSEQ or polypeptides encoded by NSEQ may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments
10 produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use. Monoclonal antibodies to PSEQ may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. In
15 addition, techniques developed for the production of chimeric antibodies can be used. (See, for example, Meyers, supra.) Alternatively, techniques described for the production of single chain antibodies may be employed. Antibody fragments which contain specific binding sites for PSEQ or the polypeptide sequences encoded by NSEQ may also be generated.

Various immunoassays may be used for screening to identify antibodies having the
20 desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art.

Yet further, an agonist of a polypeptide of PSEQ or that encoded by NSEQ may be administered to a subject to treat or prevent a cancer associated with decreased expression or
25 activity of the polypeptide.

An additional aspect of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of polypeptides of PSEQ or those encoded by NSEQ, antibodies to the polypeptides, and
30 mimetics, agonists, antagonists, or inhibitors of the polypeptides. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing

compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

5 The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

 In addition to the active ingredients, these pharmaceutical compositions may contain
10 suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

15 For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

20 A therapeutically effective dose refers to that amount of active ingredient, for example, polypeptides of PSEQ or those encoded by NSEQ, or fragments thereof, antibodies of the polypeptides, and agonists, antagonists or inhibitors of the polypeptides, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by
25 calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics.

 Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

30

EXAMPLES

It is understood that this invention is not limited to the particular methodology,

protocols, and reagents described, as these may vary. It is also understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. The examples below are provide to illustrate the subject invention and are
5 not included for the purpose of limiting the invention.

I cDNA Library Construction

The cDNA library, THYMFET02, was selected to demonstrate the construction of the cDNA libraries from which novel matrix remodeling genes were derived. The THYMFET02 cDNA library was constructed from microscopically normal thymus tissue obtained from a
10 Caucasian female fetus who died at 17 weeks gestation from anencephaly. Serology was negative; family history included tobacco abuse and gastritis.

The frozen tissue was homogenized and lysed in TRIZOL reagent (1 gm tissue/10 ml TRIZOL; Life Technologies, Rockville MD), a monoplastic solution of phenol and guanidine isothiocyanate, using a POLYTRON homogenizer (PT-3000; Brinkmann Instruments,
15 Westbury NY). After a brief incubation on ice, chloroform was added (1:5 v/v), and the lysate was centrifuged. The upper chloroform layer was removed, and the RNA was precipitated with isopropanol, resuspended in DEPC-treated water, and treated with DNase for 25 min at 37°C. The mRNA was reextracted once with acid phenol-chloroform pH 4.7 and precipitated using 0.3 M sodium acetate and 2.5 volumes ethanol. The mRNA was isolated using the
20 OLIGOTEX kit (Qiagen, Chatsworth CA) and used to construct the cDNA library.

The mRNA was handled according to the recommended protocols in the SUPERScript Plasmid system (Life Technologies). The cDNAs were fractionated on a SEPHAROSE CL4B column (Amersham Pharmacia Biotech, Piscataway NJ), and those cDNAs exceeding 400 bp were ligated into pINCY 1 plasmid (Incyte Pharmaceuticals, Palo
25 Alto CA). The plasmid was subsequently transformed into DH5 α competent cells (Life Technologies).

II Isolation and Sequencing of cDNA Clones

Plasmid DNA was released from the cells and purified using the REAL Prep 96 Plasmid kit (Qiagen). This kit enabled the simultaneous purification of 96 samples in a 96-
30 well block using multi-channel reagent dispensers. The recommended protocol was employed except for the following changes: 1) the bacteria were cultured in 1 ml of sterile Terrific Broth

(Life Technologies) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) after inoculation, the cultures were incubated for 19 hours and at the end of incubation, the cells were lysed with 0.3 ml of lysis buffer; and 3) following isopropanol precipitation, the plasmid DNA pellet was resuspended in 0.1 ml of distilled water. After the last step in the protocol, samples were
5 transferred to a 96-well block for storage at 4°C.

The cDNAs were prepared using a MICROLAB 2200 (Hamilton, Reno NV) in combination with DNA ENGINE thermal cyclers (PTC200; MJ Research, Watertown MA) and sequenced by the method of Sanger et al. (1975, J Mol Biol 94:441f) using ABI PRISM 377 DNA sequencing systems.

10 III Selection, Assembly, and Characterization of Sequences

The sequences used for coexpression analysis were assembled from EST sequences, 5' and 3' longread sequences, and full length coding sequences. Selected assembled sequences were expressed in at least three cDNA libraries.

The assembly process is described as follows. EST sequence chromatograms were
15 processed and verified. Quality scores were obtained using PHRED (Ewing et al. (1998) Genome Res 8:175-185; Ewing and Green (1998) Genome Res 8:186-194). Then the edited sequences were loaded into a relational database management system (RDBMS). The EST sequences were clustered into an initial set of bins using BLAST with a product score of 50. All clusters of two or more sequences were created as bins. The overlapping sequences
20 represented in a bin correspond to the sequence of a transcribed gene.

Assembly of the component sequences within each bin was performed using a modification of PHRAP, a publicly available program for assembling DNA fragments (Phil Green, University of Washington, Seattle WA). Bins that showed 82% identity from a local pair-wise alignment between any of the consensus sequences were merged.

25 Bins were annotated by screening the consensus sequence in each bin against public databases, such as GBpri and GenPept from NCBI. The annotation process involved a FASTn screen against the GBpri database in GenBank. Those hits with a percent identity of greater than or equal to 70% and an alignment length of greater than or equal to 100 base pairs were recorded as homolog hits. The residual unannotated sequences were screened by FASTx
30 against GenPept. Those hits with an E value of less than or equal to 10^{-8} are recorded as homolog hits.

Sequences were then reclustered using BLASTn and Cross-Match, a program for rapid protein and nucleic acid sequence comparison and database search (Green, supra), sequentially. Any BLAST alignment between a sequence and a consensus sequence with a score greater than 150 was realigned using cross-match. The sequence was added to the bin
5 whose consensus sequence gave the highest Smith-Waterman score amongst local alignments with at least 82% identity. Non-matching sequences created new bins. The assembly and consensus generation processes were performed for the new bins.

IV Coexpression Analyses of Known Matrix-remodeling Genes

Twenty one known matrix-remodeling genes were selected to identify novel genes that
10 are closely associated with matrix remodeling. The known genes were osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, II, and IV (coll-I, coll-II, and coll-III), connective tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptor (fibr-r), fibulin 1, heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor 1 (IGF 1), insulin-like growth factor binding protein
15 (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix metalloproteases (MMP), and tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3). The protein products of the known matrix-remodeling genes may be categorized as follows.

1. Extracellular matrix component protein. These proteins include collagens, proteoglycans, fibrillin, fibronectin, fibulin, and laminin that constitute the major structures of
20 the extracellular matrix.
2. Matrix proteases and matrix protease inhibitors. These proteins include matrix metalloproteases (MMPs) such as the collagenases, and MMP inhibitors such as the tissue-inhibitors of matrix metalloproteases (TIMPs).
3. Regulatory proteins that control expression of matrix-remodeling genes. Such
25 regulatory proteins include connective tissue growth factor, insulin-like growth factor, osteonectin (BM-40), and the receptors for and inhibitors of these proteins.

The known matrix-remodeling genes that we examined in this analysis, and brief descriptions of their functions, are listed in Table 4. Detailed descriptions of their roles in matrix remodeling may be found in the cited articles and reviews.

30 Table 4. Known Matrix-remodeling Genes.

Gene	Description & References
BM-40	<p>Alternate names: SPARC, osteonectin</p> <p>Regulates connective tissue remodeling, wound healing, angiogenesis</p> <p>Induces matrix metalloprotease synthesis (collagenase & gelatinase)</p> <p>Regulates cell movement and proliferation</p> <p>Expression increased in neoplastic melanoma, fibrosis, angiogenesis.</p> <p>(Kamihagi et al. (1994) <i>Biochem Biophys Res Commun</i> 200:423-8; Lane et al. (1994) <i>J Cell Biol</i> 125:929-43; Inagaki et al. (1996) <i>Life Sci</i> 58:927-34; Ledda et al. (1997) <i>J Invest Dermatol</i> 108:210-4; Shankavaram et al. (1997) <i>J Cell Physiol</i> 173:327-34.)</p>
5	
C/DSPG	<p>Chondroitin/dermatan sulfate proteoglycans</p> <p>Major extracellular matrix proteoglycan</p> <p>Regulate cell proliferation, attachment and migration.</p> <p>Darnell et al. (1990) <i>Molecular Cell Biology</i>, Scientific American Press, New York NY; Toole (1991) In: <i>Cell Biology of Extracellular Matrix</i>, Plenum, New York NY, pp. 305-341; Beck et al. (1993) <i>Biochem Biophys Res Commun</i> 190:616-23)</p>
10	
Collagens	<p>Family of fibrous structural proteins (collagen I, II, III, IV, etc.)</p> <p>Most abundant structural component of the extracellular matrix</p> <p>Secreted as procollagen; converted to collagen by MMPs</p> <p>(Alexander and Werb (1991) In: <i>Cell Biology of Extracellular Matrix</i>, pp. 255-302; Adams (1993) In: <i>Extracellular Matrix</i>, Marcel Dekker, New York, NY pp. 91-119; Schuppan et al. (1993) In: <i>Extracellular Matrix</i>, pp. 201-254.)</p>
15	
CTGF	<p>Connective tissue growth factor</p> <p>Mediates induction of matrix synthesis and fibrosis</p> <p>(Grotendorst (1997) <i>Cytokine Growth Factor Rev</i> 8:171-9; Oemar and Luscher (1997) <i>Arterioscler Thromb Vasc Biol</i> 17:1483-9; Ito et al. (1998) <i>Kidney Int</i> 53:853-61.)</p>
20	
fibrillin	<p>Major component of extracellular microfibrills (matrix elastic network)</p> <p>Present in connective tissue throughout the body</p> <p>(Kielty and Shuttleworth (1995) <i>Int J Biochem Cell Biol</i> 27:747-60; Haynes et al. (1997) <i>Br J Dermatol</i> 137:17-23; Hayward and Brock (1997) <i>Hum Mutat</i> 10:415-23.)</p>
25	
fibronectins	<p>Family of extracellular matrix glycoproteins</p> <p>Anchor cells to the matrix</p> <p>Bind matrix proteins to cell surface receptors</p>
30	
fibr-r	<p>Fibronectin receptor</p> <p>Fibronectin receptors regulate cell adhesion & migration</p> <p>(Darnell et al. (1990) <i>Molecular Cell Biology</i>, Scientific American Press, New York NY; Ruoslahti (1991) <i>Cell Biology of Extracellular Matrix</i>, pp. 343-363; Yamada (1991) <i>Cell Biology of Extracellular Matrix</i>, pp. 111-146.)</p>
fibulin 1	<p>Fibronectin-binding extracellular matrix protein</p>

- Mediates platelet adhesion via a bridge of fibrinogen
 Cleaved by matrix metalloproteinases
 Inhibits breast and ovarian cancer cell motility
 (Argraves et al. (1990) J Cell Biol 111:3155-64; Sasaki et al. (1996) Eur J Biochem 240:427-34; Hayashido et al. (1998) Int J Cancer 75:654-8.)
- 5 HSPG Heparan sulfate proteoglycans
 Extracellular matrix proteoglycan found on cell surface of many cell types
 Regulate cell interactions with the extracellular matrix
 Bind to collagens and fibronectin in the matrix
 10 Regulate cell proliferation, attachment and migration
 (Darnell et al. (1990) ; Toole (1991) In: Cell Biology of Extracellular Matrix, pp. 305-341; Schuppan et al. (1993) In: Extracellular Matrix, pp. 201-254.)
- hevin Extracellular matrix protein
 Homolog to BM-40
 15 Regulates cell adhesion and migration
 Downregulated in metastatic prostate cancer, lung cancer
 (Girard and Springer (1996) J Biol Chem 271:4511-7; Bendik et al. Cancer Res 58:232-6.)
- IGF 1 Insulin-like growth factor
 Regulates matrix homeostasis and remodeling
 20 Regulates aggregation, growth and survival of cancer cells
 (Aston et al. (1995) Am J Respir Crit Care Med 151:1597-603; Bitar and Labbad (1996) J Surg Res 61:113-9; Guvakova and Surmacz (1997) Exp Cell Res 231:149-62; Sunic et al. (1998) Endocrinology 139:2356-62.)
- IGFBP Insulin-like growth factor binding protein
 Regulates IGF-1 bioavailability (binds IGF-1 more strongly than the receptor)
 25 Degraded by matrix metalloproteases
 (Kiefer et al. (1991) Biochem Biophys Res Commun 176:219-25; Fowlkes et al. (1995) Prog Growth Factor Res 6:255-63; Parker et al. (1996) J Biol Chem 271:13523-9.)
- laminin Major protein in basal lamina, with collagen, HSPG, and entactin
 Anchors cells to the matrix by binding collagen, HSGP and heparin
 30 Laminins and collagens are the main targets of MMPs
 Regulates cell attachment, migration, growth, and differentiation
 (Yamada et al. (1993) In: Extracellular Matrix, pp. 49-66; Giannelli et al. (1997) Science 277:225-8; Quaranta and Plopper (1997) Kidney Int 51: 1441-6; Soini et al. (1997) Hum Pathol 28:220-6.)
- lumican Extracellular proteoglycan
 35 Organizes collagen fibrils in extracellular matrix
 (Dourado et al. (1996) Osteoarthritis Cartilage 4:187-96; Scott (1996) Biochemistry 35:8795-9; Cs-Szabo et al. (1997) Arthritis Rheum 40:1037-45.)
- MGP Matrix Gla protein
 Regulates calcification of cartilage

Marker for osteoblast activity

(Shanahan et al. (1994) J Clin Invest 93:2393-402; Luo et al. (1997) Nature 386:78-81; Martinetti et al. (1997) Tumour Biol 18:197-205)

- 5 MMP Family of Matrix Metalloproteases (including collagenases)
 Cleave procollagen to produce collagen
 (Alexander and Werb (1991) In: Cell Biology of Extracellular Matrix, pp. 255-302; Adams (1993) In: Extracellular Matrix, pp. 91-119; Schuppan et al. (1993) In: Extracellular Matrix pp. 201-254.)
- 10 TIMP 1, 2, 3 Tissue inhibitors of matrix metalloproteinases
 Bind and inactivate matrix proteases
 (Schuppan et al. (1993) In: Extracellular Matrix, pp. 201-254; Zvibel and Kraft (1993) In: Extracellular Matrix, pp. 559-580.)

15 The coexpression of the 21 known genes with each other is shown in Table 5. The entries in Table 5 are the negative log of the p-value ($-\log p$) for the coexpression of the two genes. As shown, the method successfully identified the strong association of the known genes among themselves, indicating that the coexpression analysis method of the present invention was effective in identifying genes that are closely associated with matrix remodeling.

Table 5. Coexpression of 21 known matrix-remodeling genes. ($-\log p$)

		laminin	fibrillin	lumican	coll IV	TIMP-1	IGFBP	coll VI	TIMP-3	CTGF	hevin	fibulin	BM-40	TIMP-2	HSPG	fibronec	MGP	C/DSP	fibr-r	coll-I	coll-III	MMP
20	laminin		7	9	21	9	15	8	4	5	7	14	10	7	11	9	19	11	7	16	10	13
	fibrillin	7		13	8	6	7	14	11	4	7	12	7	8	4	8	6	13	6	11	12	11
	lumican	9	13		24	17	16	28	17	17	14	15	22	10	8	12	25	33	14	32	34	17
	coll IV	21	8	24		17	22	22	13	11	14	28	25	12	22	16	27	26	12	34	25	26
	TIMP-1	9	6	17	17		20	15	11	11	6	10	21	15	9	16	20	13	8	14	20	19
25	IGFBP	15	7	16	22	20		20	18	16	11	14	18	14	19	21	25	23	10	27	23	20
	coll VI	8	14	28	22	15	20		13	17	19	16	20	11	11	19	19	28	12	31	36	27
	TIMP-3	4	11	17	13	11	18	13		13	18	20	22	14	9	10	18	25	12	12	13	9
	CTGF	5	4	17	11	11	16	17	13		8	10	18	7	7	19	22	12	12	18	13	11
	hevin	7	7	14	14	6	11	19	18	8		15	18	13	8	8	23	27	10	14	11	8
30	fibulin	14	12	15	28	10	14	16	20	10	15		19	9	11	8	19	20	6	17	20	18
	BM-40	10	7	22	25	21	18	20	22	18	18	19		14	11	24	21	24	16	25	32	19
	TIMP-2	7	8	10	12	15	14	11	14	7	13	9	14		7	12	8	16	11	13	13	13
	HSPG	11	4	8	22	9	19	11	9	7	8	11	11	7		8	14	10	6	11	10	10
	fibronec	9	8	12	16	16	21	19	10	19	8	8	24	12	8		14	14	11	24	21	15
35	MGP	19	6	25	27	20	25	19	18	22	23	19	21	8	14	14		32	14	25	20	13
	C/DSPG	11	13	33	26	13	23	28	25	12	27	20	24	16	10	14	32		14	27	28	14
	fibr-r	7	6	14	12	8	10	12	12	10	6	16	11	6	11	14	14		14	13	6	
	coll-I	16	11	32	34	14	27	31	12	18	14	17	25	13	11	24	25	27	14		42	21
	coll-III	10	12	34	25	20	23	36	13	13	11	20	32	13	10	21	20	28	13	42		23
40	MMP	13	11	17	26	19	20	27	9	11	8	18	19	13	10	15	13	14	6	21	23	

45 V Novel Genes Associated with Matrix Remodeling

Using coexpression analysis, we have identified 20 novel genes that show strong association with known matrix remodeling genes from a total of 41,419 assembled gene

sequences. The degree of association was measured by probability values and has a cutoff of p value less than 0.00001. This was followed by annotation and literature searches to insure that the genes that passed the probability test have strong association with known matrix-remodeling genes. This process was reiterated so that the initial 41,419 genes were reduced to the final 20 matrix-remodeling genes. Details of the coexpression patterns for the 20 novel matrix-remodeling genes are presented in Table 6.

Each of the 20 novel genes is coexpressed with at least two of the 21 known genes with a p-value of less than 10^{-7} . The coexpression results are shown in Table 6.

The novel genes identified are listed in the table by their Incyte clone numbers (Clone), and the known genes their abbreviated names (Gene) as shown in Example IV.

Table 6. Coexpression of 20 novel genes with known matrix-remodeling genes. (- log p)

Gene	laminin	fibrillin	lumican	coll IV	TIMP-1	IGFBP	coll VI	TIMP-3	CTGF	hevin	fibulin	BM-40	TIMP-2	HSPG	fibronectin	MGP	C/DSP	fibrin	coll-I	coll-III	MMP
606132	8	7	2	6	4	7	7	2	4	4	4	3	3	4	4	3	2	2	5	3	10
627722	3	4	1	1	3	3	2	5	3	6	3	4	3	2	6	5	3	3	2	3	4
639644	6	7	11	10	3	4	7	3	14	6	6	9	6	2	9	8	5	6	9	7	6
1362659	6	5	6	7	6	9	10	9	8	8	7	6	8	6	7	9	9	7	10	5	5
1446685	6	6	11	13	4	7	8	5	7	5	10	9	5	9	5	9	8	6	8	10	7
1556751	3	7	7	8	8	9	9	8	7	6	5	5	7	8	4	10	11	3	7	6	8
1656953	6	8	6	2	5	7	8	5	6	9	3	7	4	3	4	10	8	7	4	4	5
1662318	9	3	6	10	7	9	5	5	8	8	6	8	5	9	6	8	6	4	7	7	9
1996726	3	4	7	7	6	5	8	3	10	2	2	3	2	2	9	3	6	6	8	11	6
2137155	3	2	6	3	4	2	2	4	6	4	2	9	4	2	8	4	4	4	5	2	5
2268890	9	13	7	9	8	11	8	9	5	5	8	7	8	5	8	8	11	3	11	7	11
2305981	3	2	4	6	3	4	3	5	5	6	7	5	2	2	2	7	6	4	3	2	2
2457612	3	3	3	5	2	4	4	2	8	4	5	5	2	2	7	8	6	6	5	4	8
2814981	6	3	5	7	4	6	7	2	2	5	5	5	3	6	5	4	6	1	6	4	7
3089150	4	6	11	8	5	10	13	9	14	10	11	10	7	6	8	11	16	11	9	7	5
3206667	8	5	10	9	7	5	6	4	9	4	7	8	4	4	7	13	12	4	8	8	6
3284695	7	6	7	14	8	7	6	14	8	18	12	9	10	8	6	18	10	5	13	6	6
3481610	3	2	4	4	3	6	4	6	6	7	4	5	1	5	5	7	5	3	3	2	2
3722004	6	4	8	10	13	9	7	13	8	9	11	12	11	5	10	9	12	3	7	7	6
3948614	11	8	6	17	8	13	12	5	5	11	12	7	11	13	4	7	7	4	14	11	10

VI Novel Genes Associated with Matrix Remodeling

The 20 novel genes were identified from the data shown in Table 6 to be associated with matrix remodeling.

The nucleotide sequences comprising the consensus sequences of SEQ ID NOs:1-20 of the present invention were first identified from Incyte Clones 606132, 627722, 639644, 1362659, 1446685, 1556751, 1656953, 1662318, 1996726, 2137155, 2268890, 2305981, 2457612, 2814981, 3089150, 3206667, 3284695, 3481610, 3722004, and 3948614, respectively, and assembled according to Example III. BLAST and other motif searches were performed for SEQ ID NOs:1-20 according to Example VII. The sequences of SEQ ID NOs:1-20 were translated and sequence identity was sought with known sequences. Polypeptide sequences comprising the consensus sequences of SEQ ID NO:21, SEQ ID NO:22, and SEQ ID NO:23 of the present invention were encoded by SEQ ID NO:2,

SEQ ID NO:6, and SEQ ID NO:11, respectively. SEQ ID NOs:21-23 were analyzed using BLAST and other motif search tools as disclosed in Example VII.

SEQ ID NO:3 is 2987 residues in length and shows about 59% sequence identity from about nucleotide 2117 to about nucleotide 2914 with the cDNA encoding regulatory subunit of a human
5 cAMP-dependent protein kinase, RIIbeta (WO 88/03164). SEQ ID NO:8 is 3017 nucleotides in length and shows about 70% to about 74% sequence identity from about nucleotide 1 to about nucleotide 1260 and about nucleotide 1925 to about nucleotide 1985 with human Hpast mRNA (g2529706), a gene associated with multiple endocrine neoplasia type 1. SEQ ID NO:9 is 1735 nucleotides in length and shows about 25% sequence identity from about nucleotide 5 to about nucleotide 1534 with a human
10 neuronal cell adhesion molecule (WO 96/04396) important in the development of nervous system by promoting cell-cell adhesion. SEQ ID NO:14 is 2040 nucleotides in length and shows about 60% to 70% sequence identity from about nucleotide 1 to about nucleotide 1023 with a human mRNA for a serine protease (g1621243) specific for insulin-like growth factor-binding proteins. The amino acid sequence encoded by SEQ ID NO:14 from about nucleotide 3 to about nucleotide 1043 shows about
15 61% sequence identity with an osteoblast-like cell-derived protein (J09107980) useful for treatment and prevention of various diseases and as contraceptive. SEQ ID NO:15 is 2121 nucleotides in length and shows 60-80% sequence identity with a mouse gene, ADAMT-1 (g2809056), a member of the ADAM (the disintegrin and metalloproteinase) family. ADAMT-1 has been shown to contain the thrombospondin (TSP) type I motif; expression of ADAMT-1 is closely associated with inflammatory
20 processes (Kuno et al (1997) Genomics 46:466-471). SEQ ID NO:16 is 2900 nucleotides in length and shows about 70% sequence identity with a mouse homeobox (Pmx) mRNA (g460124). Homeobox genes are expressed in very specific temporal and spatial pattern and function as transcriptional regulators of developmental processes (Kern et al. (1994) Genomics 19:334-340).

SEQ ID NO:21 is 551 amino acid residues long and shows about 37% sequence identity from
25 about amino acid residue 10 to about amino acid residue 278 with PALM (g3219602), a human paralemin that is membrane-bound and expressed abundantly in brain and at intermediate levels in the kidney and in endocrine cells. In addition, the sequence encompassing residues 418 to 434 of SEQ ID NO:21 resembles one of the structural fingerprint regions of a seven trans-membrane receptor, LCR1, that is isolated from the human brain (Rimland et al. (1991) Mol Pharmacol 40:869-875). SEQ ID
30 NO:21 also has one potential amidation site at L546; three potential N-glycosylation sites at N223, N229, and N408; one potential cAMP- and cGMP-dependent protein kinase phosphorylation site at S486; fifteen potential casein kinase II phosphorylation sites at S57, S100, T101, T116, S135, S253, T349, S370, T387, S426, T434, S489, S505, S520, and T526; one potential N-myristoylation site at G54; and nine potential protein kinase C phosphorylation sites at T15, S25, S57, S100, S123, S247,

S364, S370, and S505. SEQ ID NO:22 is 99 amino acid residues in length. The sequence of SEQ ID NO:22 from about amino acid residue 71 to about amino acid residue 81 resembles one of the fingerprint regions of the RH1 and RH2 opsins, a family of G protein coupled receptors that mediate vision (Zuker et al. (1985) Cell 40:851-858; Cowman et al. (1986) Cell 44:705-710). SEQ ID NO:22 also has one potential N-myristoylation site at G24, and two potential protein kinase C phosphorylation sites at S13 and S89. SEQ ID NO:23 is 493 amino acid residues in length and shows about 44% sequence identity from about amino acid residue 277 to about amino acid residue 487 with an angiopoietin-like factor from the human cornea, CDT6 (g2765527). Angiopoietin 1 and angiopoietin 2 function as a natural ligand and a natural inhibitor, respectively, for TIE2, a receptor critical in angiogenesis during embryonic development, tumor growth, and tumor metastasis. The sequences encompassing amino acid residues 305 to 343, 346 to 355, 365 to 402, 411 to 424, and 428 to 458 of SEQ ID NO:23 resemble the carboxy-terminal domain signatures of fibrinogen beta and gamma chains from BLOCKS analysis. SEQ ID NO:23 also exhibits one potential signal peptide region encompassing amino acid residues M1 to G22 when analyzed using a HMM-based signal peptide analysis tool. In addition, SEQ ID NO:23 shows two potential N-glycosylation sites at N164 and N192; one potential cAMP- and cGMP-dependent protein kinase phosphorylation sites at S127, six potential casein kinase II phosphorylation sites at S34, S209, T238, S266, T368, and T417; four potential N-myristoylation sites at G12, G18, G22, and G29; eight potential protein kinase C phosphorylation sites at S34, S209, T268, T299, T335, S373, S383, and S477; and three potential tyrosine kinase phosphorylation sites at Y183, Y392, and Y467.

VII Homology Searching for Matrix-Remodeling Renes and the Proteins Encoded by the Genes

Polynucleotide sequences, SEQ ID NOs:1-20, and polypeptide sequences, SEQ ID NOs: 21-23, were queried against databases derived from sources such as GenBank and SwissProt. These databases, which contain previously identified and annotated sequences, were searched for regions of similarity using Basic Local Alignment Search Tool (BLAST; Altschul (1990) supra) and Smith-Waterman alignment (Smith et al. (1992) Protein Engineering 5:35-51). BLAST searched for matches and reported only those that satisfied the probability thresholds of 10^{-25} or less for nucleotide sequences and 10^{-8} or less for polypeptide sequences.

The polypeptide sequences were also analyzed for known motif patterns using MOTIFS, SPSCAN, BLIMPS, and Hidden Markov Model (HMM)-based protocols. MOTIFS (Genetics Computer Group, Madison WI) searches polypeptide sequences for patterns that match those defined in the Prosite Dictionary of Protein Sites and Patterns (Bairoch et al. supra), and displays the patterns found and their corresponding literature abstracts. SPSCAN (Genetics Computer Group) searches for

potential signal peptide sequences using a weighted matrix method (Nielsen et al. (1997) Prot Eng 10:1-6). Hits with a score of 5 or greater were considered. BLIMPS uses a weighted matrix analysis algorithm to search for sequence similarity between the polypeptide sequences and those contained in BLOCKS, a database consisting of short amino acid segments, or blocks, of 3-60 amino acids in length, compiled from the PROSITE database (Henikoff et al. supra; Bairoch et al. supra), and those in PRINTS, a protein fingerprint database based on non-redundant sequences obtained from sources such as SwissProt, GenBank, PIR, and NRL-3D (Attwood et al. (1997) J Chem Inf Comput Sci 37:417-424). For the purposes of the present invention, the BLIMPS searches reported matches with a cutoff score of 1000 or greater and a cutoff probability value of 1.0×10^{-3} . HMM-based protocols were based on a probabilistic approach and searched for consensus primary structures of gene families in the protein sequences (Eddy, supra; Sonnhammer, supra). More than 500 known protein families with cutoff scores ranging from 10 to 50 bits were selected for use in this invention.

VIII Labeling and Use of Individual Hybridization Probes

Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μCi of [γ - ^{32}P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (NEN Life Science Products, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine resin column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (NEN Life Science Products).

The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (NYTRAN PLUS, Schleicher & Schuell, Durham NH). Hybridization is carried out under the following conditions: 5x SCC/0.1% SDS at 60° C for about 6 hours, subsequent washes are performed at higher stringency with buffers, such as 1x SCC/0.1% SDS at 45° C, then 0.1xSCC. After XOMAT AR film (Eastman Kodak, Rochester NY) is exposed to the blots for several hours, hybridization patterns are compared.

IX Production of Specific Antibodies

SEQ ID NO:20, 21, or 23 substantially purified using polyacrylamide gel electrophoresis (Harrington (1990) Methods Enzymol 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the amino acid sequence is analyzed using LASERGENE software (DNASTAR, Madison WI) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for

selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. Typically, oligopeptides 15 residues in length are synthesized using an ABI 431A peptide synthesizer (PE Biosystems) using Fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester to increase

- 5 immunogenicity. Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for anti-peptide activity by, for example, binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

10

What is claimed is:

1. A substantially purified polynucleotide comprising a gene that is coexpressed with one or
5 more known matrix-remodeling genes in a plurality of biological samples, wherein each known
matrix-remodeling gene is selected from the group consisting of osteonectin, chondroitin/dermatan
sulfate proteoglycans, collagen I, II, II, and IV, connective tissue growth factor, fibrillin, fibronectins,
fibronectin receptor, fibulin 1, heparan sulfate proteoglycan, extracellular matrix protein, insulin-like
growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein,
10 matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3.

2. The polynucleotide of claim 1, comprising a polynucleotide sequence selected from the
group consisting of:

- (a) a polynucleotide sequence selected from the group consisting of SEQ ID NOs:1- 20;
- (b) a polynucleotide sequence which encodes the polypeptide sequence of SEQ ID NO: 21,
15 22, or 23;
- (c) a polynucleotide sequence having at least 70% identity to the polynucleotide sequence of
(a) or (b);
- (d) a polynucleotide sequence comprising at least 18 sequential nucleotides of the
polynucleotide sequence of (a), (b), or (c);
- 20 (e) a polynucleotide which hybridizes under stringent conditions to the polynucleotide of (a),
(b), (c), or (d); and
- (f) a polynucleotide sequence which is complementary to the polynucleotide sequence of (a),
(b), (c), (d), or (e).

3. A substantially purified polypeptide comprising the gene product of a gene that is
25 coexpressed with one or more known matrix-remodeling genes in a plurality of biological samples,
wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin,
chondroitin/dermatan sulfate proteoglycans, collagen I, II, II, and IV, connective tissue growth factor,
fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular
matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin,
30 lumican, matrix Gla protein, matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase
1, 2, and 3.

4. The polypeptide of claim 3, comprising a polypeptide sequence selected from the group
consisting of:

- (a) the polypeptide sequence of SEQ ID NO:21, 22, or 23;
- 35 (b) a polypeptide sequence having at least 85% identity to the polypeptide sequence of (a);
and

(c) a polypeptide sequence comprising at least 6 sequential amino acids of the polypeptide sequence of (a) or (b).

5. An expression vector comprising the polynucleotide of claim 2.

6. A host cell comprising the expression vector of claim 5.

5 7. A pharmaceutical composition comprising the polynucleotide of claim 2 or the polypeptide of claim 3 in conjunction with a suitable pharmaceutical carrier.

8. An antibody which specifically binds to the polypeptide of claim 4.

9. A method for diagnosing a disease or condition associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes, wherein each known
10 matrix-remodeling gene is selected from the group consisting of osteonectin, chondroitin/dermatan sulfate proteoglycans, collagen I, II, III, and IV, connective tissue growth factor, fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein, matrix metalloproteinases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3, the method
15 comprising the steps of:

(a) providing a sample comprising one or more of said coexpressed genes;

(b) hybridizing the polynucleotide of claim 2(F) to said coexpressed genes under conditions effective to form one or more hybridization complexes; and

(c) detecting the hybridization complexes, wherein the altered level of hybridization
20 complexes compared with the level of hybridization complexes of a nondiseased sample correlates with the presence of the disease or condition.

10. A method for treating or preventing a disease associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a subject in need, wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin,
25 chondroitin/dermatan sulfate proteoglycans, collagen I, II, III, and IV, connective tissue growth factor, fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein, matrix metalloproteinases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3, the method comprising the step of administering to said subject in need the pharmaceutical
30 composition of claim 7 in an amount effective for treating or preventing said disease.

11. A method for treating or preventing a disease associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a subject in need, wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin, chondroitin/dermatan sulfate proteoglycans, collagen I, II, III, and IV, connective tissue growth factor,

fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein, matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3, the method comprising the step of administering to said subject in need the antibody of
5 claim 8 in an amount effective for treating or preventing said disease.

12. A method for treating or preventing a disease associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a subject in need, wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin, chondroitin/dermatan sulfate proteoglycans, collagen I, II, III, and IV, connective tissue growth factor,
10 fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein, matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3, the method comprising the step of administering to said subject in need the polynucleotide sequence of claim 2(F) in an amount effective for treating or preventing said disease.

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.
WALKER, Michael G.
VOLKMUTH, Wayne
KLINGLER, Tod M.

<120> MATRIX-REMODELING GENES

<130> PB-0004 PCT

<140> To Be Assigned

<141> Herewith

<150> 09/169,289

<151> 1998-10-09

<160> 23

<170> PERL Program

<210> 1

<211> 1447

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> 1380

<223> a or g or c or t, unknown, or other

<220>

<221> misc_feature

<223> Incyte ID No.: 606132CB1

<400> 1

```

cctggaacca gaaggagacc tacctgcaca tcatgaagaa cgaggaggag gtggtgatct 60
tgttcgcgca ggtgggcgac cgcagcatca tgcaaagcca gagcctgatg ctggagctgc 120
gagagcagga ccaggtgtgg gtacgcctct acaagggcga acgtgagaac gccatcttca 180
gcgaggagct ggacacctac atcaccttca gtggctacct ggtcaagcac gccaccgagc 240
cctagctggc cggccacctc ctttcctctc gccaccttcc acccctgcgc tgtgctgacc 300
ccaccgcctc ttccccgatc cctggactcc gactccctgg ctttggcatt cagtgaagacg 360
ccctgcacac acagaaaagcc aaagcgatcg gtgctcccag atcccgacgc ctctggagag 420
agctgacggc agatgaaatc accagggcgg ggcacccgcg agaaccctct gggaccttcc 480
gcggccctct ctgcacacat cctcaagtga ccccgcacgg cgagacgcgg gtggcggcag 540
ggcgtcccag ggtgcggcac cgcgggtcca gtccttgaa ataattaggc aaattctaaa 600
ggtctcaaaa ggagcaaagt aaaccgtgga ggacaaagaa aagggttgtt atttttgtct 660
ttccagccag cctgctggct cccaagagag aggccttttc agttgagact ctgcttaaga 720
gaagatccaa agttaaaagt ctggggtcag gggaggggcc gggggcagga aactacctct 780
ggcttaattc ttttaagcca cgtaggaact ttcttgagg ataggtggac cctgacatcc 840
ctgtggcctt gcccaagggc tctgctggtc tttctgagtc acagctgcga ggtgatgggg 900
gctggggccc caggcgctcag ctcccagagg gacagctgag cccctgcct tggctccagg 960
ttggtagaag cagccgaagg gctcctgaca gtggccaggg acccctgggt cccccaggcc 1020
tgcagatgtt tctatgaggg gcagagctcc tggtacatcc atgtgtggct ctgctccacc 1080
cctgtgccac cccagagccc tggggggtgg tctccatgcc tgccaccctg gcacggcctt 1140
tctgtgccgc ctcccacaca aatcagcccc agaaggcccc ggggccttgg cttctgtttt 1200
ttataaaaca cctcaagcag cactgcagtc tcccatctcc tcgtgggcta agcatcaccg 1260
cttcacagtg tgttgtgttg gttggcagca aggctgatcc agacccttc tgccccact 1320
gcgctcatcc aggcctctga ccagtagcct gagaggggct ttttctaggc ttcagagcan 1380
gggagagctg gacggggtag acagtccgct tgtctgttct aagctctgtg agctcagctt 1440
gagacaa                                     1447

```

<210> 2
<211> 2481
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No.: 627722CB1

<400> 2
ctagcaagca ggtaaacgag ctttgtacaa acacacacag accaacacat ccggggatgg 60
ctgtgtgttg ctagagcaga ggctgattaa acactcagtg tgttggtctt ctgtgccact 120
cctggaaaat aatgaattgg gtaaggaaaca gttaataaga aaatgtgcct tgctaactgt 180
gcacattaca acaaaagagct ggcagctcct gaaggaaaag ggcttgtgcc gctgccgttc 240
aaacttgtca gtcaactcat gccagcagcc tcagcgtctg cctccccagc acaccctcat 300
tacatgtgtc tgtctggcct gatctgtgca tctgctcgga gacgctcctg acaagtctgg 360
aatttctcta tttctccact ggtgcaaaga gcggatttct ccttcttctt cttctgtcac 420
ccccgctcct ctcccccagg aggtccttg atttatggta gctttggact tgcttccccg 480
tctgactgtc cttgacttct agaatggaag aagctgagct ggtgaaggga agactccagg 540
ccatcacaga taaaagaaaa atacaggaag aaatctcaca gaagcgtctg aaaatagagg 600
aagacaaaat aaagcaccag catttgaaga aaaaggcctt gagggagaaa tggcttctag 660
atggaatcag cagcggaaaa gaacaggaag agatgaagaa gcaaaatcaa caagaccag 720
accagatcca ggttctagaa caaagtatcc tcaggcttga gaaagagatc caagatcttg 780
aaaaagctga actgcaaadc tcaacgaagg aagaggccat tttaaagaaa ctaaagtcaa 840
ttgagcggac aacagaagac attataagat ctgtgaaagt ggaaagagaa gaaagagcag 900
aagagtcaat tgaggacatc tatgctaata tccctgacct tccaaagtc tacatacctt 960
ctagggttaag gaaggagata aatgaagaaa aagaagatga tgaacaaaat aggaaagctt 1020
tatatgccat ggaaattaaa gttgaaaaag acttgaagac tggagaaaagt acagttctgt 1080
cttcaatacc tctgccatca gatgacttta aaggtacagg aataaaagtt tatgatgatg 1140
ggcaaaagtc agtgtatgca gtaagttcta atcacagtg agcatacaat ggcaccgatg 1200
gcctggcacc agttgaagta gaggaacttc taagacaagc ctacagagaga aactctaat 1260
ccccaacaga gtatcatgag cctgtatatg ccaatccctt ttacaggcctt acaacccccac 1320
agagagaaac ggtgacccct ggaccaaact ttcaagaaag gataaaagatt aaaactaatg 1380
gactgggtat tgggtgaaat gaatccatac acaatatggg caatggctct tcagaggaaa 1440
ggggaaacaa cttcaatcac atcagtccca ttccgccagt gcctcatccc cgatcagtga 1500
ttcaacaagc agaagagaag cttcacaccc cgcaaaaaag gctaattgact ccttgggaag 1560
aatcgaatgt catgcaggac aaagatgcac cctctccaaa gccaaaggct agccccagag 1620
agacaatatt tgggaaatct gaacaccaga attcttcacc cacttgtcag gaggacgagg 1680
aagatgtcag atataatatc gttcattccc tgcctccaga cataaatgat acagaaccgg 1740
tgacaatgat tttcatgggg tatcagcagg cagaagacag tgaagaagat aagaagtttc 1800
tgacaggata tgatgggatc atccatgctg agctgggtgt gattgatgat gaggaggagg 1860
aggatgaagg agaagcagag aaaccgtcct accaccccat agctcccat agtcagggtgt 1920
accagccagc caaaccaaca ccacttccta gaaaaagatc agaagctagt cctcatgaaa 1980
acacaaatca taaatcccc cacaataatt ccatatctct gaaagagcaa gaagaaagct 2040
taggcagccc tgtccaccat tccccatttg atgctcagac aactggagat gggactgagg 2100
atccatcctt aacagcttta aggatgagaa tggcaaagct gggaaaaaag gtgatctaag 2160
agttgtacca cctatataaa catcctttga agaagaaact aagaagcatt tgcaaatctt 2220
tcttctggat attttgttta tttttctga agtccaaaaa attatcatta cagtgtacca 2280
tattaagcca tgtgaataag tagtagtcat tatttgtgaa aaattcccaa aaagctgggg 2340
aaaacaaatg tgtaactttt ccagttactt gacacgattc agtgggggaa aaccagcatt 2400
ttttattcta ttgataccaa agcatttcta ataagagctt gttaaattta agaataaagt 2460
tatttaaaat aaaaaaaaaa a 2481

<210> 3
<211> 2987
<212> DNA
<213> Homo sapiens

<220>
<221> unsure

<222> 2955

<223> a or g or c or t, unknown, or other

<220>

<221> misc_feature

<223> Incyte ID No.: 639644CB1

<400> 3

agaaaaaaag	aaaaaagaaa	aaaactaagg	cagcagctct	taataaataa	cacctggagc	60
agaatcggta	aactgctttc	acgttggctt	ttgcagaagt	ggcaatgcat	tgaggatata	120
tctggcaagc	ttcgaattca	caagtgtaaa	ggacccagtg	acctgctcac	agtccggcag	180
agcacgcgga	acctctacgc	tcgcggcttc	catgacaaag	acaaagagtg	cagttgtagg	240
gagtctgggt	accgtgccag	cagaagccaa	agaaagagtc	aacggcaatt	cttgagaaac	300
caggggactc	caaagtacaa	gccagattt	gtccatactc	ggcagacacg	ttccttgtcc	360
gtcgaatttg	aaggtgaaat	atatgacata	aatctggaag	aagaagaaga	attgcaagtg	420
ttgcaaccaa	gaaacattgc	taagcgctcat	gatgaaggcc	acaaggggcc	aagagatctc	480
caggcttcca	gtggtggcaa	caggggcagg	atgctggcag	atagcagcaa	cgccgtgggc	540
ccacctacca	ctgtccgagt	gacacacaag	tgttttattc	ttcccaatga	ctctatccat	600
tgtgagagag	aactgtacca	atcggccaga	gcgtggaagg	accataaggc	atacattgac	660
aaagagattg	aagctctgca	agataaaatt	aagaatttaa	gagaagttag	aggacatctg	720
aagagaagga	agcctgagga	atgtagctgc	agtaaacaaa	gctattacaa	taaagagaaa	780
ggtgtaaaaa	agcaagagaa	attaaagagc	catcttcacc	cattcaagga	ggctgctcag	840
gaagtagata	gcaaactgca	acttttcaag	gagaacaacc	gtaggaggaa	gaaggagagg	900
aaggagaaga	gacggcagag	gaagggggaa	gagtgcagcc	tgcttggcct	cacttgcttc	960
acgcatgaca	acaaccactg	gcagacagcc	ccgttctgga	acctgggac	tttctgtgct	1020
tgacagagtt	ctaacaataa	cacctactgg	tgtttgcgta	cagttaatga	gacgcataat	1080
tttcttttct	gtgagtttgc	tactggcttt	ttggagtatt	ttgatatgaa	tacagatcct	1140
tatcagctca	caaatacagt	gcacacggta	gaacgaggca	ttttgaatca	gctacacgta	1200
caactaatgg	agctcagaag	ctgtcaagga	tataagcagt	gcaacccaag	acctaagaat	1260
cttgatgttg	gaaataaaga	tggaggaagc	tatgacctac	acagaggaca	gttatgggat	1320
ggatgggaag	gttaatcagc	cccgtctcac	tgacagacac	aactggcaag	gcctagagga	1380
gctacacagt	gtgaatgaaa	acatctatga	gtacagacaa	aactacagac	ttagtctggt	1440
ggactggact	aattacttga	aggattttaga	tagagtattt	gcactgctga	agagtctcag	1500
tgagcaaaat	aaaacaaaata	agactcaaac	tgtcaaagt	gacgggttct	tggttgtctc	1560
tgctgagcac	gctgtgtcaa	tggagatggc	ctctgctgac	tcagatgaag	acccaaggca	1620
taaggttggg	aaaacacctc	atttgacctt	gccagctgac	cttcaaacc	tgcatattga	1680
ccgaccaaca	ttaagtccag	agagtaaact	tgaatggaat	aacgacattc	cagaagttaa	1740
tcatttgaat	tctgaacact	ggagaaaaac	cgaaaaatgg	acggggcatg	aagagactaa	1800
tcactctgaa	accgatttca	gtggcgtatg	catgacagag	ctagagctcg	ggcccagccc	1860
caggctcgag	cccattcgca	ggcaccgcaa	agaacttccc	cagtatggtg	gtcctggaaa	1920
ggacattttt	gaagatcaac	tatatcttcc	tgtgcattcc	gatggaattt	cagttcatca	1980
gatgttcacc	atggccaccg	cagaacaccg	aagtaattcc	agcatagcgg	ggaagatggt	2040
gaccaagggt	gagaagaatc	acgaaaagga	gaagtccacg	cacctagaag	gcagcgccct	2100
ctcttcactc	tcctctgatt	agatgaaact	gttaccttac	cctaaacaca	gtatttcttt	2160
ttaacttttt	tatttgtaaa	ctaataaagg	taatcacagc	caccaacatt	ccaagctacc	2220
ctgggtacct	ttgtgcagta	gaagctagtg	agcatgtgag	caagcgggtg	gcacacggag	2280
actcatcggt	ataatttact	atctgccaa	agtagaaaga	aaggctgggg	atatttgggt	2340
tggcttgggt	ttgatttttt	gcttgtttgt	ttgttttgta	ctaaaacagt	attatctttt	2400
gaatatcgta	gggacataag	tatatcatg	ttatccaatc	aagatggcta	gaatggtgcc	2460
tttctgagtg	tctaaaactt	gacacccctg	gtaaatcttt	caacacactt	ccactgcctg	2520
cgtaatgaag	ttttgattca	tttttaacca	ctggaatttt	tcaatgccgt	cattttcagt	2580
tagatgattt	tgcaatttga	gattaaaatg	ccatgtctat	ttgattagtc	ttattttttt	2640
attttttacag	gcttatcagt	ctcactgttg	gctgtcattg	tgacaaagtc	aaataaaacc	2700
ccaaggacga	cacacagtat	ggatcacata	ttgttttgaca	ttaagctttt	gccagaaaat	2760
gttgcatgtg	ttttacctcg	acttgctaaa	atcgattagc	agaaaggcat	ggctaataat	2820
gttggtgggt	aaaataaata	aataagtaaa	caaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	2880
aaaaaaaaaa	aaaaaaaaaa	aaaaagcaaa	aaaagctgcc	gccacagtta	gatgaagaag	2940
catgaggatc	cgagnnggtc	gcctctttga	gtggtgaggg	agtcgcg		2987

<210> 4

<211> 2915

<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<223> Incyte ID No.: 1362659CB1

<400> 4

```
gaggcaagaa ttccggcacga gggacatttt gccaaacttaa acgagaaaaa gacccccgc 60
accgggcaca ctcccccttc ctccagcccc gcttcagcca catgctccag ctgctgcca 120
gtaaagccct gtgccttttt ttccccgaa tactgcccac agcatccctt tcccatctgc 180
ctctcaggag ttggggactt tgctaggaga ttttttaagt gttccttact gggacaacgt 240
ggagccacgt ttgcaggagc tccatttgta tccctgctgg tgttgacttc tgtgtagggg 300
ccagttcatg tccctgactc tcacctccca ttagataaat gaagcccacc cccctttcta 360
gagtgatgag agtcaagaag aggggatgta tgaacggcca aattcccatg tgagaggaag 420
atgacctgat ccacctagcc ttttcttctg gatctgtcct cctcacccc tttcacctga 480
gctgtccaca gtaggaaca taaagaaaca atgtccccta catatcccca tgactacata 540
atccatcatc gtaggaata ggaaagcaaa tttgattttg gttttgtaaa acgtacatgc 600
ttcaataatt ctttttttgt gtcttaataa ctcatagggg aaaaaaacag ctcacccaag 660
gtgttaggtt tcacatatat attcatcaac tatttttaga gatttaattc tatcaaactc 720
tgtattacct cagatcattt taaatagcaa gccaaataac agctttgaag gctattttac 780
cattcctgtt cacaaaaggt tctcatgggt cctgacaggt tacccttgag ggcttgtgtc 840
tactttttta aagtcaatgg tttttttct tgtgttctag ttcccataat aggagagaaa 900
atatagaaat atatgcaaaa attatagttt tctttagatc agaaactgat atttttgggt 960
cagccatatg tattttgttt aaaggattta aaataaagtg ccgtcatgta gccctgtgga 1020
agggagcaca taaccagctg tttggcatga caggtgactt agtatatttg taattgggtt 1080
taaaaccaat acaccatact ttctttctgc aaacagccat ctttatactt agggaagaaa 1140
aattgttggg ttctagactt ttttaataa aattttgttg atatggaatt aggttaagttt 1200
aagtgtctat gtgcataatg tttttatata agttttttct attcagtttc actgatccaa 1260
ctggcagtggt gtaaataatg cataagttta taacactttt ccccaaatg gtgctttgga 1320
tttgaaaagg gtctgatggg gagaaggaga acgtatcatc ctagcttctt ctcttaataa 1380
acctagaaaa acgggtagta aactgtggat agtcaggaaa acaccagca agggacacag 1440
ctgtcaggaa atgaatcttc cccccaaccc ccaccatgca gatggataga cagaatcttt 1500
cctgactagt cattagatc aggggcctct gttggatttg tgtttcttga agaatagtctg 1560
gcagagtggg ataaaagaca cgaatatctc ctggtctata aggatactct gatttggggg 1620
ttgcattttt catggttttt atttcctgtt cccctggag tttccatta gtgagttttt 1680
gtgcaaggat cttatttgtg atgccttccc tcccctagaa agattttgtg caatatatta 1740
aatggggaca gaattctaaa tggataaaac aatggctggg tctagccctg agtgacagtc 1800
ttaaggctag atccttccca tagtatcatc tgtcctctgg aatgactctc ctgtccctaa 1860
aggggttaag agagagatca cctagaaatc cctctggaca cttgtgggtt ctttaggggt 1920
tgagtttctt cttccccttg agcttcagag aggagagttg gcatggttta atctgaatgg 1980
ttacctact gctgaaaacc cagagggcg tggcacactc gcttgtgtgg aaaagcctct 2040
aaatgcatcc ctctcttctt ttctgtcttc ctttgctta caattgaagc agcccggtgt 2100
accatcacag tatgcagaga ctctctcacc tttcatatct agggaccacc cccgatgcat 2160
tggtgagggt gggcacttat aaatgcctgc tattgttaag ccattccagc ctcttctct 2220
gaatagacca gacgcccttt cacttagttc agtgccagtc cttttgcctt cccaacctgt 2280
ctgttaggcc tgctgttccc tttgctcttg attaggagag atggaaggag atgagctccc 2340
ataactgaat tggccttttg ttcatgtttt ctcccatat gtatatatgc catatgtgaa 2400
tatgccatat atatgtgcc acaaatctat ctacgttggt cttttcaaat tagcacgcag 2460
ataggaattt tgagtttctt cttcttttag taactagtat aacaagcact ggtatttttg 2520
tacaaaaaag aaaaacaaaa gattgactat tgtggtctgc atgacataaa caaacaatg 2580
gtgatataca agcaacgtat accccagtc agtgtgtgtt gccataattt gcaattcagc 2640
ttaacagtgc acccaatcta tatttgcatt ttgatattat ttaagctcta tgtacaagg 2700
tttgcagtga tttatatggt tcttagggaa aaaaaatgct ataaactgca aatctgaaat 2760
tcaaatgtgt tgttccactg agaccagaag aagaagagga gttttaaaag ggataatttg 2820
ttggagccaa taaagctttt tgctgatgaa cagaaaccaa tactgctgtg cactgagaat 2880
aaaaactcat gccacttgt aaaaaaaaaa aaagg 2915
```

<210> 5

<211> 1826
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No.: 1446685CB1

<400> 5
gaaagcgcca gcctcagtcg cgccgcccgc cgtgcgtcc gccagcgcc agctccgcgt 60
cccagccggc ccgcccgcgc ctgcgccgcg ccatggccac ctcccgcag aagtcgcctt 120
ctgtcccaa gtctcccact cccaagtcgc ccccgctccg caagaaagat gattccttct 180
tggggaaact cggaggggacc ctggcccggg ggaagaaagc caaggagggtg tccgagctgc 240
aggaggaggg aatgaacgcc atcaacctgc ccctcagccc aattcccttt gagctggacc 300
ccgaggacac gatgctggag gagaatgagg tgcgaacaat ggtggatcca aactcacgca 360
gtgaccccaa gcttcaagaa ctgatgaagg tattaattga ctggattaat gatgtgttgg 420
ttggagaaa aatcattgtg aaagacctag ctgaagattt gtatgatgga caagtcctgc 480
agaagctttt cgagaaactg gagagtgaga agctaaatgt ggctgagggt acccagtcag 540
agattgtctc gaagcaaaaa ctgcagactg tcctggagaa gatcaatgaa accctgaaac 600
ttcctcccag gagcatcaag tggaatgtgg attctgttca tgccaagagc ctggtggcca 660
tcttacacct gctcgttgct ctgtctcagt atttccgcgc accaattcga ctccagacc 720
atgtttccat ccaagtgtt gtggtccaga aacgagaagg aatcctccag tctcggcaaa 780
tccaagagga aataactggt aacacagagg ctctttccgg gaggcataaa cgtgatgcct 840
ttgacacctt gttcgaccat gcccagaca agctgaatgt ggtgaaaaag aactcatca 900
ctttcgtgaa caagcacctg aataaactga acctggagggt cacagaactg gaaacccagt 960
ttgcagatgg ggtgtacctg gtgctgctca tggggctcct ggagggtac tttgtgccc 1020
tgcacagctt cttcctgacc ccggacagct ttgaacagaa ggtcttgaat gtctcctttg 1080
cctttgagct catgcaagat ggagggttg aaaagccaaa accgcggcca gaagacatag 1140
tcaactgtga cctgaaatct aactacagag tgtgtacaa cctcttcacc aagtaccgta 1200
acgtggagtg aggggctgcc ctgggcccac cactgcccac gagttccttg tgttggcgta 1260
ctggaccctc ctccgaactg cttaccctg cttattcctg tctcttgca tgtgctctcc 1320
cacaagtcca gctgcaaccc agagatagt gaaactgaaa ttaggaagga aatcatcaat 1380
aactcagtg gctgaccat ccctcccagg cgctggggac caacctagca atgaagggtg 1440
ggaaagttgt tcccttcccgt gtgccagggt cagatttccc tccatgattt ggggaaccagg 1500
ttaggcaaaa gagtccccac aagatgaaaa taaagatcct agttaccatt caaaggatgc 1560
taactgtgtg tcaggcccca cactaagtgc tctgctctga tatactcaag gccattaatc 1620
ttcaggactc ccattgacgt aggtgtttca ttcccctttt acagatgagg aaactaaggc 1680
ttggagggtt aatgacttgc cagaagttgg aatttttttc ctctttgaac ataacctctc 1740
ccttctccct aaaggtaacc actattctga gtccaatcat caaggttttg cttttctttt 1800
tagctaagta tcaattcctc aatagt 1826

<210> 6
<211> 1439
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No.: 1556751CB1

<400> 6
gagtatccct tgtttaatca cttttgtggt taaaagagac ctttgggtca gtctgcctca 60
ttccttgaag agtttagccc tggctcactt ttcactctat ttcttctcct gtctcaagaa 120
agaagaaaaa aagagacaaa ttaccagaa acccctccct tcccacatg gaggccttgg 180
caaatgttaa ttttcttaga aaatccttca gacctgaaga cgcaggaaaa gaatctggct 240
ctcagggtgg cttctgcgtc cccgcccga gggcccagac tatggtcaca gggcgtcct 300
gttctctccc gggactccag aatttctctc ctcaaaggaa agaaaaacagg gcatgcgctt 360
gttggaacaa cgcaggggccg gctcccaaaa accccatgtg tgtacgatta aaagttggcc 420
gtccccaggc ctcccagcgc aaacttaaa agacagggtt ttgctgaaaa ccaaactatg 480
gccagctggg ctttttaaca acctagagac tttccggagc tgcttgaac agagcctgcg 540
ggaaacgggg cttgccagag aactcacag tttccttcat ggctgtttt ggtcccttaa 600
gaatctccac atcattgtct ttcttgtgcc ttttcttgg tgagcaacag aaagggaagg 660


```
gttccaagcc tctaaaaatg tgctttgtga tcaggagtgc gctccaaacc aaatacgcg 720
gctgcccttt cgaggccagt gagctcagcc tccaaggctt taaagccaca tttcagcaag 780
agaaagcgct gagagctcgc aggttcatta aagaaggcaa agcactggtt tctctcctta 840
gaaaagtagg tttcttggct tgatgtagac tggcttgctt tgatttttag tgaaggggaat 900
gtacgtaaaa caaaataggg cttggctggt caaaggagac aagcaggatg gatggatgga 960
tggatggatg gatgtatgga tgaatagata gatggtggtt gcatgtaaat tgcagagaaa 1020
acaaaaccaa agctgattgg aaacaattaa ttgtgggtgt ctgaggggga aggtcgcgagc 1080
tttgggcagc tttgagaagc ggtacaagag ttctgtgctt gtgtgtccag ccctggagcc 1140
agccagtgca tttattttta gctcttagaa gcaactcctt ggcccaggaa tgcgtgaccc 1200
ctgagatggg tccacgcac tctctacact tcttctctc cgtgggatac tggactcgtg 1260
cctctgcgcc cattctcttc tcacgcatac ccatgagctt taatttctact ttctgatcac 1320
ggtacgtcca taaagccagt attacactta aatgaagtat tcttttttgt aatcgttttt 1380
tttagaaggt aaacaaattt aataaagcta ccaataatga gaaaaaaaaa aaaaaaaaaa 1439
```

<210> 7

<211> 3047

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 1656953CB1

<400> 7

```
cgagacagag gaaatgtgtc tccctccaag gccccaaagc ctcagagaaa ggggtgtttct 60
ggttttgcct tagcaatgca tcggtctctg aggtgacact ctggagcggg tgaagggcca 120
caaggtgcag ggtaataact cttgccagtt ttgaaatata gatgctatgg ttcagattgt 180
ttttaataga aaactaaagg ggcaggggaa gtgaaaggaa agatggaggt tttgtgcggc 240
tcgatggggc atttggaact tctttttaaa gtcactcat ggtctccagt tttcagttgg 300
aactctgggt tttaacactt aaggagagaca aaggctgtgt ccatttggca aaacttcctt 360
ggccacgaga cttcaggtga tgtgtgaagc ttggcagctc gtggtgtgga gacgagccat 420
gtgtctggcc attcagagga ttctaaagac atggctggat gcgctgctga ccaacatcag 480
cacttaataa aatgcaaatg caacatttct cctctgggc cttgaaaatc cttgccctta 540
tcatttgggg tgaaggagac atttctgtcc ttggcttccc acagcccaa cgcagtctgt 600
gtatgattcc tgggatccaa cgagccctcc tattttcaca gtgttctgat tgctctcaca 660
gcccaggccc atcgtctgtt ctctgaatgc agccctgttc tcaacaacag ggaggtcatg 720
gaaccctct gtggaaccca caaggggaga aatgggtgat aaagaatcca gttcctcaaa 780
accttccctg cgaggtggg tccctctcct gctgggtggt gcttctctt gcacaccact 840
cccaccacgg ggggagagcc agcaacccaa ccagacagct caggttgtgc atctgatgga 900
aaccactggg ctcaaacacg tgctttattc tctgtttat ttttgctgtt actttgaagc 960
atggaaattc ttgtttgggg gatcttgggg ctacagtagt gggtaaacaa atgccaccg 1020
gccaagaggc cattaacaaa tcgtccttgt cctgaggggc cccagcttgc tcgggcgtgg 1080
cacagtgggg aatccaaggg tcacagtagt gggagaggtg caccctgcca cctgctaact 1140
tctcgctaga cacagtgtt ctgccaggt gacctgttca gcagcagaac aagccagggc 1200
catggggacg ggggaagttt tcacttggag atggacacca agacaatgaa gatttgttgt 1260
ccaaataggc caataattct gggagactct tggaaaaaac tgaatatatt caggaccaac 1320
tctctccctc cctcatccc acatctcaaa gcagacaatg taaagagaga acatctcaca 1380
caccagctc gccatgccta ctcatctctg aatttcaggt gccatcactg ctctttcttt 1440
cttctttgtc atttgagaaa ggatgcagga ggacaattcc cacagataat ctgaggaatg 1500
cagaaaaacc agggcaggac agttatcgac aatgcattag aacttgggtg gcatcctctg 1560
tagagggact ccacccctgc tcaacagctt ggcttccagg caagaccaac cacatctggt 1620
ctctgccttc ggtggccac acacctaagc gtcacgtca ttgccatagc atcatgatgc 1680
aacacatcta cgtgtagcac tacgacgtta tgtttgggta atgtggggat gaactgcatg 1740
aggctctgat taaggatgtg gggaagtggg ctgcggtcac tgtcggcctt gcaaggccac 1800
ctggaggcct gtctgttagc cagtgggtga ggagcaaggc ttcaggaagg gccagccaca 1860
tgccatcttc cctgcgatca ggcaaaaaag tggaaataaa aagtcaaaacc ttatatgca 1920
tgtgttatgt ccattttgca ggatgaactg agtttaaaag aatttttttt tctcttcaag 1980
ttgctttgtc ttttccatcc tcatcacaag cccttgtttg agtgtcttat ccctgagcaa 2040
tctttcgatg gatggagatg atcattaggt acttttgttt caacctttat tcctgtaaat 2100
atttctgtga aaactaggag aacagagatg agatttgaca aaaaaaaatt gaattaaaaa 2160
taacacagtc tttttaaaac taacatagga aagcctttcc tattatttct cttcttagct 2220
tctcattgtg ctacattcagg aaaacaggaa aacacagctt tctagcagct gcaaaatggt 2280
```

ttaatgcccc	ctacatat	ccatcacctt	gaacaatagc	tttagcttgg	gaatctgaga	2340
tatgatccca	gaaaacatct	gtctctactt	cggctgcaaa	acccatgggt	taaactctata	2400
tggtttgtgc	attttctcaa	ctaaaaatag	agatgataat	ccgaattctc	catatatattca	2460
ctaatacaag	acactat	catactagat	tcctgagaca	aatactcact	gaagggcttg	2520
tttaaaaaata	aattgtgttt	tggctctgttc	ttgtagataa	tgcccttcta	ttttaggttag	2580
aagctctgga	atccctttat	tgtgctgttg	ctcttatctg	caaggtggca	agcagttctt	2640
ttcagcagat	tttgcccaact	attcctctga	gctgaagttc	tttgcataga	tttggtctaa	2700
gcttgaatta	gatccctgca	aaggcttgct	ctgtgatgtc	agatgtaatt	gtaaatgtca	2760
gtaatcactt	catgaacgct	aaatgagaat	gtaagtattt	ttaaatgtgt	gtatttcaaaa	2820
tttgtttgac	taattctgga	attacaagat	ttctatgcag	gatttacctt	catcctgtgc	2880
atgtttccca	aactgtgagg	agggaaggct	cagagatcga	gcttctcctc	tgagttctaa	2940
caaaatgggtg	ctttgaggggt	cagccttttag	gaaggtgcag	ctttgttgctc	ctttgagctt	3000
tctgttatgt	gcctatccta	ataaactctt	aaacacaaaa	aaaaaaa		3047

<210> 8

<211> 3017

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 1662318CB1

<400> 8

cgcaaactca	accctttcgg	aaacaccttc	ctcaacaggt	tcattgtgtgc	ccagctccct	60
aatcagggtc	tggagagcat	cagcatcctc	gacaccccg	gtatcctgtc	gggtgccaag	120
cagagagtga	gccgcggcta	cgacttccc	gccgtgctgc	gctgggttcg	ggagcgcgtg	180
gacctcatca	tcctgtctct	tgatgcgcac	aagctggaga	tctcggacga	gttctcagag	240
gccatcggcg	cgttgccggg	ccatgaggac	aagatccgcg	tgggtgctcaa	caaggccgac	300
atgggtggaga	cgcagcagct	gatgcgcgtc	tacggcgcgc	tcattgtggg	gctgggcaag	360
gtgggtgggca	cgcccgaggt	gctgcgcgtc	tacatcgggt	ccttctgggt	ccagcccctc	420
ctgggtgccc	acaaccggcg	cctcttcgag	ctggaggagc	aggacctctt	ccgcgacatc	480
cagggcctgc	cccggcacgc	agccttgccg	aagctcaacg	acctgggtgaa	gagggcccg	540
ctgggtgcgag	ttcacgctta	catcatcagc	tacctgaaga	aggagatgcc	ctctgtgttt	600
gggaaggaga	acaagaagaa	gcagctgate	ctcaaactgc	ccgtcatctt	tgcaagatt	660
cagctggaac	atcacatctc	ccctggggac	tttccctgatt	gccagaaaat	gcaggagctg	720
ctgatggcgc	acgacttcac	caagtttccac	tcgctgaagc	cgaagctgct	ggaggcactg	780
gacgagatgc	tgacgcacga	catcgccaag	ctcatgcccc	tgctgcggca	ggaggagctg	840
gagagcaccc	aggtggggcgt	gcaggggggc	gcttttgagg	gcacccacat	gggcccgttt	900
gtggagcggg	gacctgacga	ggccatggag	gacggcgagg	agggctcgga	cgacgaggcc	960
gagtgggtgg	tgaccaagga	caagtccaaa	tacgacgaga	tcttctacaa	cctggcgcc	1020
gccgacggca	agctgagcgg	ctccaaggcc	aagacctgga	tgggtggggac	caagctcccc	1080
aactcagtgc	tggggcgcct	ctggaagctc	agcgtatgtg	accgcgacgg	catgctggat	1140
gatgaagagt	tcgcgctggc	cagccacctc	atcgaggcca	agctgggaag	ccacgggctg	1200
cccgcccaac	tgccccgtcg	cctgggtgcc	ccctccaaag	gacgccacaa	gggctccgcc	1260
gagtgaagcc	ggcccccttc	ccatggccct	gctgtggctc	cccagctcca	gtcggctgca	1320
cgcacacccc	tgctccggct	cacacacgcc	ctgcctgccc	tccttgcccc	gctgtaagga	1380
ccgggggtct	ccctcctcac	taccgccaga	caccccggtg	gaagcattta	gaggggacca	1440
cgggagggac	aaggcttctc	tgctccgcct	tcacacctcc	agcctcacgt	tcacttaggc	1500
acatcacaca	cacactggca	cacgcaggca	tcctccatc	cgctattcat	tcaaatattt	1560
attgagcacc	tactatgtgc	ccagccctgt	tctaggcaat	gggcattacc	atagagaaca	1620
aaatagacaa	atacatctgc	cctcatggaa	ggtgacgttc	ccaggagagg	gcacctacac	1680
agtcacgcaa	acacacacta	attcctggca	gggccccag	ccctccctct	ggctgagcag	1740
ccctgtggct	gaaatgacta	gcagataaac	agacccctt	ctgctccgct	tcctcctgcc	1800
cagccaggca	acaccctcaa	ccggctccat	cacatcctca	ggtctcggga	ccatgggggg	1860
ctcagagggg	agacacacct	actgcttctc	cagatgggcc	cctccgcagc	cccttccctt	1920
gctcgggaa	agcccccaat	tctgcccaca	ccatttatt	tccttctctc	cttctctctt	1980
ttctttcctt	ccttctctct	ttttgtttt	tgcccccaat	tctgcccata	ccatttctt	2040
tccttctctc	cttctctctt	ttttgtttt	gccccagtt	ctgtccacac	cccttccctt	2100
tcctgtcctg	tcctttcttt	cttttttgat	agaatcttgc	tctgtcgccc	aggctgggag	2160
tgcaagtggg	agatctcagc	tcaactgcaac	ctccacctcc	tgggttgaag	tgattctcgt	2220
gcctcagcct	cctgagtagc	tgggactgca	ggcacgcgcc	accacgcccc	gctaattttt	2280

gtattttgagt	agagacgggg	tttcaccatg	ttggccaggc	tggtctcgaa	ctccgcacat	2340
caggtgatct	gctcgcctcg	gcctcccaaa	gtgatgggat	tacaggcatg	agccaccgtg	2400
cccggcttca	cacccatttc	tttaaaaagg	atcccgtagc	aggcagaaaa	gccccctcca	2460
tcctgctcct	ctgatactgt	gcccccttgg	agatatttcc	gtcctccacc	cacgtgtctg	2520
tggttggaac	tgcccagcct	gctcctggcc	ccctggaagc	ctccccacag	ctggtaatct	2580
ggactttaagg	attgctgggc	caccgcctct	ctgcctacca	ccattccata	tttaagtggg	2640
gccccctacgt	agaaaaggccc	cggggcttta	ttttagtctc	cttttcaggg	atgtcgtggg	2700
cgggggaggg	ggttcttggg	gctacagccc	tctccccacc	cctaaaggga	cgccgacgct	2760
gtttgctgcc	ttcaccacat	attagtgcct	gaccctggca	ggggacccca	tggaaaagat	2820
ggggaagagc	aaaatacatg	gagacgacgc	accctccagg	atgctcgtctg	ggattcccac	2880
gccccaccat	gtccccccacc	ccatggctgg	gaggggcctc	tgaacggaac	agtgtcccca	2940
cagagcgaat	aaagcaaggc	ttcttcccca	aaaaaaaaaa	aaaaaaaaaa	attggtgcgg	3000
ccgaagtatt	tcccttc					3017

<210> 9

<211> 1735

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 1996726CB1

<400> 9

tcgggaggaa	ggagactaca	cctgctttgc	tgaaaatcag	gtcgggaagg	acgagatgag	60
agtcagagtc	aaggtggtga	cagcgcccg	caccatccgg	aacaagactt	acttggcggt	120
tcaggtgccc	tatggagacg	tggtcactgt	agcctgtgag	gccaaaggag	aacccatgcc	180
caaggtgact	tggttgctcc	caaccaacaa	ggtgatcccc	acctcctctg	agaagtatca	240
gatataccaa	gatggcactc	tccttattca	gaaagcccag	cggtctgaca	gcggcaacta	300
cacctgcttg	gtcaggaaca	gcgcgggaga	ggtaggaag	acggtgtgga	ttcacgtcaa	360
cgtccagcca	cccaagatca	acggtaaccc	caacccccatc	accaccgtgc	gggagatagc	420
agccgggggc	agtcggaaac	tgattgactg	caaagctgaa	ggcatcccca	ccccgagggt	480
gttatgggct	tttcccagag	gtgtgggtct	gccagctcca	tactatggaa	accggatcac	540
tgtccatggc	aacggttccc	tggacatcag	gagtttgagg	aagagcgact	ccgtccagct	600
ggtatgcatg	gcacgcaacg	agggagggga	ggccagggtg	atcgtgcagc	tcactgtcct	660
ggagcccatg	gagaaaccca	tcttcacaga	cccgatcagc	gagaagatca	cggccatggc	720
gggcccacacc	atcagcctca	actgctctgc	cgcggggacc	ccgacaccca	gcctggtgtg	780
ggtccttccc	aatggcaccg	atctgcagag	tggacagcag	ctgcagcgct	tctaccacaa	840
ggctgacggc	atgctacaca	ttagcgggtct	ctcctcgggtg	gacgccgggg	cctaccgctg	900
cgtggcccg	aatgccgctg	gccacacgga	gaggctggtc	tccctgaagg	tgggactgaa	960
gccagaagca	aacaagcagt	atcataacct	ggtcagcatc	atcaatgggtg	agaccctgaa	1020
gctccccctgc	acccctcccg	gggctgggca	gggacgtttc	tcctggacgc	tccccaatgg	1080
catgcatctg	gagggccccc	aaaccttggg	acgcgtttct	cttctggaca	atggcaccct	1140
cacggttcgt	gaggcctcgg	tgtttgacag	gggtacctat	gtatgcagga	tggagacgga	1200
atacggccct	tcggtaacca	gcatccccgt	gattgtgatc	gcctatcctc	cccggatcac	1260
cagcgagccc	accccggtca	tctacacccg	gcccgggaac	accgtgaaac	tgaactgcat	1320
ggctatgggg	attcccaaag	ctgacatcac	gtgggagtta	ccggataagt	cgcactctgaa	1380
ggcaggggtt	caggctcgtc	tgtatggaaa	cagatttctt	cacccccagg	gatcactgac	1440
catccagcat	gccacacaga	gagatgccgg	cttctacaag	tgcattggcaa	aaaacattct	1500
cggcagtgc	tccaaaacaa	cttacatcca	cgtcttctga	aatgtggatt	ccagaatgat	1560
tgcttaggaa	ctgacaacaa	agcgggggtt	gtaagggaag	ccaggttggg	gaataggagc	1620
tcttaaataa	tgtgtcacag	tgcattgggtg	cctctgggtg	gtttcaagtt	gaggttgatc	1680
ttgatctaca	attgtttggg	aaaggaagca	atgcagacac	gagaaggagg	gctca	1735

<210> 10

<211> 1016

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 2137155CB1

<400> 10

ctgtacgttc	ccctgtggcc	cacgcctagt	gaaaatgata	tcgtacatct	ccctagagat	60
atgggtcacc	tccaggtaga	ttacagagat	aacaggctgc	acccaagtga	agattcttca	120
ctggactcca	ttgcctcagt	tgtggttccc	ataattatat	gcctctctat	tataatagca	180
ttcctattca	tcaatcagaa	gaaacagtgg	ataccactgc	tttgcctgga	tcgaacacca	240
actaagcctt	cttccttaaa	taatcagcta	gtatctgtgg	actgcaagaa	aggaaccaga	300
gtccaggttg	acagttccca	gagaatgcta	agaattgcag	aaccagatgc	aagattcagt	360
ggcttctaca	gcatgcaaaa	acagaacctat	ctacaggcag	acaatttcta	ccaaacagtg	420
tgaagaaagg	caactaggat	gaggtttcaa	aagacggaag	acgactaaat	ctgctctaaa	480
aagtaaacta	gaatttgctg	acttgcttag	tggattgtat	tggattgtga	cttgatgtac	540
agcgctaaga	ccttactggg	atgggctctg	tctacagcaa	tgtgcagaac	aagcattccc	600
acttttcctc	aagataactg	accaagtgtt	tcttagaacc	aaagttttta	aagttgctaa	660
gatatatttg	cctgtaagat	agctgtagag	atatttgggg	tggggacagt	gagtttggat	720
ggcgaaatac	accgcacggg	ggtgttggga	agaaaaattt	gtcagcttgg	ctcggggaga	780
aaccctggta	cactaaagca	gttcagtgtg	ccagagggtta	tttttttccc	attgctctga	840
agactgcact	ggttgctgca	aactcaggcc	tgaatgagcg	gaaacaaaaa	aagccttgcg	900
ccccgatgcc	ataacacctt	tggaatcccc	agcgccctc	agaaaccttt	tcaggcatcc	960
aggtcttaag	cccaagtatc	tttctataca	gtcccactgc	ggtgagcgtg	ggggag	1016

<210> 11

<211> 2288

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 2268890CB1

<400> 11

caaccagggt	caggctgtgc	tcacagtttc	ctctggcgcc	atgtaaaggc	tccacaaagg	60
agtgtggagt	tcaaagtagg	ctgctgcgga	cggcctgagg	atggacccca	agccctggac	120
ctgccgagcg	tggcactgag	gcagcggtcg	acgctactgt	gagggaaaga	aggttgtgag	180
cagccccgca	ggacccctgg	ccagccctgg	ccccagcctc	tgccggagcc	ctctgtggag	240
gcagagccag	tggagcccag	tgaggcaggg	ctgcttggca	gccaccggcc	tgcaactcag	300
gaacccctcc	agaggccatg	gacaggctgc	cccgtgacg	gccagggtga	agcatgtgag	360
gagccgcccc	ggagccaagc	aggagggaa	aggctttcat	agattctatt	cacaaagaat	420
aaccaccatt	ttgcaaggac	catgaggcca	ctgtgcgtga	catgctgggtg	gctcggactg	480
ctggctgcc	tgggagctgt	tgacggccag	gaggacggtt	ttgagggcac	tgaggagggc	540
tcgccaaagag	agttcattta	cctaaacagg	tacaagcggg	cgggcgagtc	ccaggacaag	600
tgcacctaca	ccttcattgt	gccccagcag	cgggtcacgg	gtgccatctg	cgtcaactcc	660
aaggagcctg	aggtgcttct	ggagaaccga	gtgcataagc	aggagctaga	gctgctcaac	720
aatgagctgc	tcaagcagaa	gcggcagatc	gtagcgtgc	agcagctggt	ggaggtggac	780
ggcgcgattg	tgagcgaggt	gaagctgctg	cgcaaggaga	gccgcaacat	gaactcgcgg	840
gtcacgcagc	tctacatgca	gctcctgcac	gagatcatcc	gcaagcggga	caacgcgttg	900
gagctctccc	agctggagaa	caggatcctg	aaccagacag	ccgacatgct	gcagctggcc	960
agcaagtaca	aggacctgga	gcacaagtac	cagcacctgg	ccacactggc	ccacaaccaa	1020
tcagagatca	tcgcgcagct	tgaggagcac	tgccagaggg	tgccctcggc	caggcccgtc	1080
ccccagccac	ccccgctgc	ccgccccgg	gtctaccaac	cacccaccta	caaccgcatc	1140
atcaaccaga	tctctaccaa	cgagatccag	agtgaccaga	acctgaaggt	gctgccaccc	1200
cctctgcccc	ctatgccac	tctcaccagc	ctcccatctt	ccaccgacaa	gccgtcgggc	1260
ccatggagag	actgcctgca	ggccctggag	gatggccacg	acaccagctc	catctacctg	1320
gtgaagccgg	agaacaccaa	ccgcctcatg	cagggtgtgt	gcgaccagag	acacgacccc	1380
gggggctgga	ccgtcatcca	gagacgcctg	gatggctctg	ttaacttctt	caggaactgg	1440
gagacgtaca	agcaagggtt	tgggaacatt	gatggcgaat	actggctggg	cctggagaac	1500
atttactggc	tgacgaacca	aggcaactac	aaactcctgg	tgacctgga	ggactgggtc	1560
ggccgcaaag	tctttgcaga	atacgccagt	ttccgcctgg	aacctgagag	cgagtattat	1620
aagctgcggc	tggggcgcta	ccatggcaat	gcgggtgact	cctttacatg	gcacaacggc	1680
aagcagttca	ccaccctgga	cagagatcat	gatgtctaca	caggaaactg	tgccactac	1740
cagaagggag	gctggtggta	taacgcctgt	gcccactcca	acctcaacgg	ggtctggtac	1800
cgcggggggc	attaccggag	ccgctaccag	gacggagtct	actgggctga	gttccgagga	1860

ggctcttact	cactcaagaa	agtgggtgatg	atgatccgac	cgaaccccaa	caccttccac	1920
taagccagct	ccccctcctg	acctctcgtg	gccattgcca	ggagcccacc	ctgggtcacgc	1980
tggccacagc	acaaagaaca	actcctcacc	agttcatcct	gaggctggga	ggaccgggat	2040
gctggattct	gttttccgaa	gtcactgcag	cggatgatgg	aactgaatcg	atacgggtgtt	2100
ttctgtccct	cctacttttc	ttcacaccag	acagcccctc	atgtctccag	gacaggacag	2160
gactacagac	aactctttct	ttaaataaat	taagtctcta	caataaaaac	acaactgcaa	2220
agtaccttca	taatatatac	gtgtatgagc	ctcccctgtg	cacgtatgtg	tatagcacat	2280
atatatgg						2288

<210> 12

<211> 3304

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 2305981CB1

<400> 12

ccctcttatg	gattcccagc	aagcatcagg	aaccattgtg	caaattgtca	tcaataacaa	60
acacaagcat	ggacaagtgt	gtgtttccaa	tggaaagacc	tattctcatg	gcgagtcctg	120
gcacccaaac	ctccgggcat	ttggcattgt	ggagtgtgtg	ctatgtactt	gtaatgtcac	180
caagcaagag	tgtaaagaaa	tccactgccc	caatcgatac	ccctgcaagt	atcctcaaaa	240
aatagacgga	aagtgtctga	aggtgtgtcc	aggtaaaaaa	gcaaaaagaag	aacttccagg	300
ccaaagcttt	gacaataaag	gtactttctg	cggggaagaa	acgatgcctg	tgtatgagtc	360
tgtattcatg	gaggatgggg	agacaaccag	aaaaatagca	ctggagactg	agagaccacc	420
tcaggtagag	gtccacgttt	ggactattcg	aaagggcatt	ctccagcact	tccatattga	480
gaagatctcc	aagaggatgt	ttgaggagct	tcctcacttc	aagctggtga	ccagaacaac	540
cctgagccag	tggaaatctt	tcaccgaagg	agaagctcag	atcagccaga	tgtgttcaag	600
tcgtgtatgc	agaacagagc	ttgaagattt	agtcaagggt	ttgtacctgg	agagatctga	660
aaagggccac	tgttaggcaa	gacagacagt	attggatagg	gtaaagcaag	aaaactcaag	720
ctgcagctgg	actgcaggct	tattttgctt	aagtcaacag	tgccctaaaa	ctccaaactc	780
aaatgcagtc	aattattcac	gccatgcaca	gcataatttg	ctcctttgtg	tggagtgggtg	840
tgtcagccct	tgaacatctc	ctccaaagag	actagaagag	tcttaaatga	tatgtgggag	900
gaggagggat	agaacatcac	aacactgctc	tagtttcttg	gagaatcaca	tttctttaca	960
gggttaaagac	aaacaagacc	ccagggtttt	tatctagaaa	gttattcaag	tgaaagaaag	1020
agaagggaat	tgcttagtag	gagttctgca	gtatagaaca	attacttgta	tgaaattata	1080
cctttgaatt	ttagaatgtc	atgtgttctt	ttaaaaaaat	tagctcccca	tcctccctcc	1140
tcactccctc	cctccctcct	tctctctctc	tctctctctc	cctctctcac	agacacacac	1200
acacacacac	acacacgcac	acgcacgtcc	acactcacat	taaactaaag	ctttatttga	1260
agcaaagcta	gccaaaattc	tacgttactt	ttcccttgac	tggatcccaa	gtagcttgga	1320
agttttttgtg	cccaggagag	taaataactg	tgaacaagag	gctctgccct	taggtccttg	1380
tggctgttta	agtcaccaac	aatagagtca	gggtaaagaa	taaaaaactt	ttcatagcct	1440
cattcattca	cttagaagtg	gtaataattt	ttccctaattg	ataccacttt	tcttttcccc	1500
ctgtacctat	gggacttcca	gaaagaagtt	aaattgagta	aaatcatcag	aaactgaatc	1560
catgtaagaa	aaaataattg	ttgaagaaag	aagttgatag	aattcaaaaa	ggccatcttt	1620
ttgctttcac	atcaataaaa	tttaccaggt	aatagatcag	tactcactaa	tattttttgag	1680
accatagttg	tctggtcaga	aaaattatat	taaattagta	aattctagaa	gctctttaa	1740
agggaaagttt	tcttcttctt	ccaattatag	gagttgattt	ttactttgca	aagtggctcg	1800
gtcctcatga	gcatctgcat	gttgactctt	cagttaagaa	aattgttggt	catttaggga	1860
gggtggatatt	ctgatgaaga	tctttatcct	aaaccttcct	actatccttg	tcttattcat	1920
caagcagata	tttttagtcaa	gaattccaga	gaaggctgct	cctaaaaatgt	ctacttgacg	1980
cccaatacca	gagcataaac	tatccattct	ggggtctggc	tttagaaatc	atcctttgtg	2040
gaagacctaa	ttcttcacag	caaggatctc	aggcatgcct	tctagatttg	ttccctctga	2100
ggggcaggaa	tgaactgtag	aaatgtttta	aggaccctga	aaccccatat	gtctctttcc	2160
atgactatag	gtgagagaat	tctttcctaa	gagggtttga	taccaatagg	ggaaaatgta	2220
aaatgtttcag	tctttatgac	aacctggcat	aaaggagtca	attcttatga	aagagacaca	2280
agggcccttat	ggccagggtt	tcttgggaca	agactctcac	cagcacatca	cacacgttct	2340
ccttggaaga	gagaagcagt	acatcccggg	tgagaggtca	caaagcatta	gtttgtgtgt	2400
gtgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	gtggtaaagg	ggggaagggtg	2460
ttatgaggct	gtccctcccg	tcccagaggt	ggcagtgatt	ccataatgtg	gagactagta	2520
actagatcct	aaggcaaaaga	ggtgtttctc	cttctggatg	attcatccca	aagccttccc	2580

```

accaggtgt tctctgaaag cttagcctta agagaacacg cagagagttt ccctagatat 2640
actcctgcct ccaggtgctg ggacacacct ttgcaaaatg ctgtgggaag caggagctgg 2700
ggagctgtgt taagtcaaag tagaaacctt ccagtgtttg gtgttggtga gagaatagga 2760
catagggtaa agaggccaag ctgcctgtag ttagtagaga agaattggatg tggttcttct 2820
tgtgtatttta tttgtatcat aaacacttgg aacaacaaag accataagca tcatttagca 2880
gtttagacca ttttctagtt aactcatgta aacaagtaag agtaacataa cagtattacc 2940
ctttcactgt tctcacagga catgtacctt attatggtag ttatttatgt agtcaactga 3000
tttctggatt tttaaattaa taaaaaagt aattttgaaa aaaaaaaaaa aaaaaaaaaa 3060
aaaaaaaaaa aaaaaaaaaa actcgagggg gggcctgtac cgggttcccc gtaacaggtt 3120
cgcccttaag attccctggc cgcagttttt ggccgcgttt tggggaacct ctgggtacct 3180
cccttagttgc tcgctaaat cccctttcgc agcccgttta aaggctgggg ccggccgatt 3240
gccttcccaa tagcctccca tgaatgggaa tggaattgga agggaaattt tggtaaattc 3300
ggtg                                     3304

```

<210> 13
 <211> 708
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No.: 2457612CB1

```

<400> 13
ggaaagccag gaagtgcagg aatcatttca tcagggccaa taactacacc acccctgagg 60
tcaacaccca ggcctactgg aactcccttg gagagaatag agacagatgt aaagcaacca 120
acagttcctg cctctggaga agaactggaa aatataactg acttttagctc aagcccaaca 180
agagaaactg atcctcttgg gaagccaaga ttcaaaggac ctcatgtgctg atacatccaa 240
aagcctgaca acagtccctg ctccattact gactctgtca aacggttccc caaagaggag 300
gccacagagg ggaatgccac cagcccacca cagaacccac ccaccaacct cactgtggtc 360
accgtggaag ggtgcccctt catttgtcat ctggactgg gaaaagccac taaatgacac 420
tgtcactgaa tatgaagtta tatccagaga aaatgggtca ttcagtggga agaacaagtc 480
cattcaaatg acaaatcaga cattttccac agtagaaaat ctgaaaccaa acacgagtta 540
tgaattccag gtgaaaccca aaaacccgct tgggtgaaggc ccggtcagca acacagtggtc 600
attcagtact gaatcagcgg acccagagtg agtgagcagt ttctgcagga gagatgcctc 660
tggtactgaag gccgctttgt tcgactcttg ctcagggtga agggcaac 708

```

<210> 14
 <211> 2040
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No.: 2814981CB1

```

<400> 14
cggccagccg ccgcgcgctg cagctctccg ggacgcccgt gcgccagctg cagaaggggc 60
cctgcccgtt ggggtctccac cagctgagca gccgcgctta caagttcaac ttcatgtctg 120
acgtggtgga gaagatcgca ccagccgtgg tccacataga gctcttcctg agacaccgc 180
tgtttggccg caacgtgccc ctgtccagcg gttctggctt catcatgtca gaggccggcc 240
tgatcatcac caatgccac gtggtgtcca gcaacagtgc tgccccgggc aggcagcagc 300
tcaaggtgca gctacagaat ggggactcct atgaggccac catcaaagac atcgacaaga 360
agtcggacat tgccaccatc aagatccatc ccaagaaaaa gctccctgtg ttgttgctgg 420
gtcactcgcg cgactgcgg cctggggagt ttgtggtggc catcggcagt cccttcgccc 480
tacagaacac agtgacaacg ggcatcgta gcactgcccc gcgggagggc agggagctgg 540
gcctccggga ctccgacatg gactacatcc agacggatgc catcatcaac tacgggaact 600
ccgggggacc actggtgaac ctggatggcg aggtcattgg catcaacacg ctcaagggtc 660
cggctggcat ctcccttgcc atccctcag accgcatcac acggttcctc acagagttcc 720
aagacaagca gatcaaagac tgaagaagc gcttcacggt catacggatg cggacgatca 780

```

```

caccaagcct ggtggatgag ctgaaggcca gcaaccgga cttcccagag gtcagcagt 840
gaatttatgt gcaagaggtt gcgccgaatt caccttctca gagaggcggc atccaagatg 900
gtgacatcat cgtcaaggtc aacgggcgtc ctctagtga ctcgagtga ctgcaggagg 960
ccgtgctgac cgagtctcct ctccactgg aggtgcggcg ggggaacgac gacctcctct 1020
tcagcatcgc acctgaggtg gtcagtgtgag gggcgcatc ctccagcgcc aagcgtcaga 1080
gcctgcagac aacggagggc agcgccccc cgagatcagg acgaaggacc accgtcggtc 1140
ctcagcaggg cggcagcctc ctctggctg tccggggcag agcggaggct gggcttggcc 1200
agggggccga atttccgcct ggggagtgtt ggatccacat cccggtgccg gggagggaag 1260
cccaacatcc ccttgtacag atgatcctga aagtcacttc caagttctcc ggatattcac 1320
aaaactgcct tccatggagg tcccctcctc tcctagcttc ccgcctctgc ccctgtgaac 1380
acccatctgc agtatccctt gtcctgccc ctctactgc aggtctgggc tgccaagctt 1440
tctcccccct gacaaacgcc cacctgacct gaggccccag ctccctctg ccctaggact 1500
taccaagctg tagggccagg gctgctgcct gccagcctgg ggtccctgga ggacagggtc 1560
catctgatcc ctttgggtg cgggggtggg gtccagccca gagcaggcac tgagtgaatg 1620
ccccctggct gcggagctga gccccgcct gccatgaggt tttcctccc aggcaggcag 1680
gagggcgcgg ggagcacgtg gaaagtggc tgctgcctgg ggaagcttct cctccccaag 1740
gcggccatgg ggcagcctgc agaggacagt ggacgtggag ctgcggggtg tgaggactga 1800
gccggcttcc ccttcccacg cagctctggg atgcagcagc cgctcgcag gaagtggccg 1860
ccagaggcat gcaggctgct gggcaccacc ccctcatcca gggaacgagt gtgtctcaag 1920
gggcatttgt gagctttgct gtaaatggat tcccagtgtt gcttgactg tatgtttctc 1980
tactgtatgg aaaataaagt ttacaagcac aaaaaaaaaa aaaaaaaaaa aaaaaaagg 2040

```

<210> 15

<211> 2121

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 3089150CB1

<400> 15

```

gtaaaagctg gttgtgatcg catcatagac tccaaaaaga agtttgataa atgtggtgtt 60
tgccggggaa atggatctac ttgtaaaaaa atatcaggat cagttactag tgcaaacct 120
ggatatcatg atatcatcac aattccaact ggagccacca acatcgaagt gaaacagcgg 180
aaccagagg gatccaggaa caatggcagc tttcttgcca tcaaagctgc tgatggcaca 240
tatattctta atggtgacta cactttgtcc accttagagc aagacattat gtacaaaggt 300
gttgtcttga ggtacagcgg ctctctgctg gcattggaaa gaattcgag ctttagccct 360
ctcaaagagc ccttgaccat ccaggttctt actgtgggca atgcccttcg acctaaaatt 420
aaatacacct acttcgtaaa gaagaagaag gaatctttca atgctatccc cactttttca 480
gcatgggtca ttgaagagtg gggcgaatgt tctaagtcac gtgaattggg ttggcagaga 540
agactggtag aatgccgaga cattaatgga cagcctgctt ccgagtgtgc aaagggaagt 600
aagccagcca gcaccagacc ttgtgcagac catccctgcc ccagtgaggc gctgggggag 660
tggtcatcat gttctaagac ctgtgggaag gggtacaaaa aaagaagctt gaagtgtctg 720
tcccatgatg gaggggtgtt atctcatgag agctgtgatc ctttaaagaa acctaaacat 780
ttcatagact tttgcacaat ggcagaatgc agttaagtgg tttaaagtgg gtttagctttg 840
agggcaaggc aaagtgagga agggctgggt cagggaaagc aagaaggctg gagggatcca 900
gcgtatcttg ccagtaacca gtgaggtgta tcagtaaggt gggattatgg gggtagatag 960
aaaaggagtt gaatcatcag agtaaaactgc cagttgcaaa tttgatagga tagttagtga 1020
ggattattaa cctctgagca gtgatatagc ataataaagc cccgggcatt attattatta 1080
tttcttttgt tacatctatt acaagtttag aaaaaacaaa gcaattgtca aaaaaagtta 1140
gaactattac aaccctgttt tcttggtact tatcaaatac ttagtatcat ggggggttggg 1200
aaatgaaaag taggagaaaa gtgagatttt actaagacct gttttacttt acctcactaa 1260
caatgggggg agaaaggagt acaaatagga tctttgacca gcactgttta tggctgctat 1320
ggtttcagag aatgtttata cattatttct accgagaatt aaaacttcag attgttcaac 1380
atgagagaaa ggctcagcaa cgtgaaataa cgcaaatggc ttccctcttc cttttttgga 1440
ccatctcagt ctttatttgt gtaattcatt ttgaggaaaa aacaactcca tgtatttatt 1500
caagtgcatt aaagtctaca atggaaaaaa agcagtgaag cattagatgc tggtaaaagc 1560
tagaggagac acaatgagct tagtacctcc aacttccttt ctttcctacc atgtaaccct 1620
gctttgggaa tatggatgta aagaagtaac ttgtgtctca tgaaaatcag tacaatcaca 1680
caaggaggat gaaacgccgg aacaaaaatg aggtgtgtag aacagggtcc cacagggtttg 1740
gggacattga gatcatttgt cttgtggtgg ggaggctgct gaggggtagc aggtccatct 1800

```

```

ccagcagctg gtccaacagt cgtatcctgg tgaatgtctg ttcagctctt ctgtgagaat 1860
atgatttttt ccatatgtat atagtaaaat atgttactat aaattacatg tactttataa 1920
gtattggttt ggggtgtcct tccaagaagg actatagtta gtaataaatg cctataataa 1980
catattttatt tttatacatt tattttctaatt gaaaaaaact tttaaattat atcgcttttg 2040
tggaagtgca tataaaaatag agtattttata caatatatgt tactagaaat aaaagaacac 2100
ttttggaaaa aaaaaaaaaa a 2121

```

```

<210> 16
<211> 2900
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No.: 3206667CB1

```

```

<400> 16
gaagttttta aaaaaactac agcagccaaa gaaactatat atatatatat atatatccag 60
aatgattgcc tctactgtcc tcattgactt gtttgaacct tagtgcctta ccctgtcctc 120
ttcccagttc tctttataga agctctagga gctttcgaaa agccaaagtc tttctgaaga 180
atctgtgctg gacagacata attccccttc tcattgtctc catctttgtt ggtcatggta 240
aggttttttc atcagcctct gaaaaaatag ttgtgcacaa catctgtctc ctggactgtc 300
tgatccaatg taattggctg cgtctggcta attctaagca cttaaagtcta catctaagct 360
atagatttaa gcttgaagct acagattata tcaactatcac caccacccct caccagtgta 420
aatcagacag tcagtcactt taagttaaag atatttgttg tctttgaatg atttgctgtc 480
acagactatt tggtagaaga aatatttttc acctgagaga ggaagagaaa tttctctagt 540
aacacaaaaga gtgagttcta aaaggcatgc ccacatctct ttcgtgcctt aaggatagtg 600
agatgcacac ttatatatat actgtatata tttatatatt tatatatata tttcatatat 660
atatataata ttgcaagctt aagtttgcaa tttcccaaac aatacaaaaa gcaaattaca 720
caccctcacc actgttctta tctctatagt gatgaaacat taattagggg tcttgctgct 780
tttctttttc tacacgaagt tttcattaaa gccacagaat aattgatagg gcagctgttt 840
gagaacaggt cccatttttc cattagggtt ttaaataaat tagaaactat ttgaggtcat 900
aaaaatgtcc ttgagtttgg agcctgagct ctggtgaaat gctgatacat ctgatctatc 960
atgggaattg cagttagaga gagtaaggaa taccatttag tcatctatcc gttcttcact 1020
tagcaggaat atgaaagaaa ggcacatgtt taagaggaat acctaaaggt ttttctaaat 1080
tccaacattt aaaaggcaat tgtgggctat ttttattttt taatattttg aaataaagtt 1140
tagtgtctag ggctgggagc caggactgat cttccatttc tttttctttg ttcccagcca 1200
tgcttttgta acttgccagg tggacttgac caactacatt accatgctgt gcctcgtgaa 1260
acccatttgt aaaatgggat taataatact tacctacctc acaggggtgt tgtgaggtc 1320
tattcatttg ctcttttatt ctttctctgt tctctgtat gtccagcact ttgtagccat 1380
gggaggaag ggactataaa agtgtacaat gttaatggaa tgatacggta cctgaaagcc 1440
ttgttttcta gtaagaaaat gctaccttgc tgtacatact tataaccttg tatttgaaa 1500
tgagaaatag gtttatattt tcagatctct caaaaatcac atcatttgac caaagataa 1560
tttaagacac atagaacaga ttttttttct ttatatattc atoctgacca gcttagttct 1620
aataattttt agttgtgagt gattaaaaaa ctttggatca attttgggtc aacatgccaa 1680
ctttgtagtc tgagtgcacg gcaaggattt ttgggtttta gatgcacttt tagcacacat 1740
ttgtatttcc cttggcatat cagattgagc taatgggtgat gttatttcaa tctaacagcc 1800
accaatctga aattgtattt caaatgttga ttctgtagtt ctttaaataa taatgaagct 1860
catcttatac attttgcttt caccaattga ttcttcttc ttttagccca ctattaaaac 1920
atttcttact gaatggttca tgtaggcttg ctgaacagca cgcattactt gcttcttgaa 1980
gagttccccc attcatccat ttgtcccat agttgctgtg gattatcaag ttttgaagga 2040
actgtacatc ccaacagact gaaacattct aagtgaatg agtataatcc aagtaactgg 2100
tgaactttgg aggtttggag cttgaagaga atggctaaga agatttgaat tatagggagg 2160
gaacagaaat catacatgaa aaggttttac tgagaagggg aaaaccttag atagagggac 2220
atgtgaaaca aaatcatttg aaattttgat tcagacatcc atttccagtg gcaaacagca 2280
aagcctgaac ccataaacc aaatgatagg tgaagtggg tggttttatc caatgtctca 2340
agcaagcaat atctgggaat atcatagagt acaagtgct ggctagccaa agaaacattc 2400
actgctggtg aaccaatacc ataagcatgt attatctaag cacttgatca agaaatatac 2460
atgttgtaac agctctcaat tttgttcatt tattatcaaa tttttaaaat acaagtttgg 2520
tatgtgattt ggaaaagatg ctttctggat ctttaagccag ttgtcagtg aggtcctcag 2580
ggctgcaaat gtcaagacat aacctgttc ctcaccatca tgataccaga tacaggtgaa 2640
tacataggaa ctatctgcct gtgtcctcaa tctcccttca aacaagatgc tgattttag 2700

```



```

gggtacttggc aggtttaaatt aaaccagaag aggtgactta ataaaaaagg gaatgacatt 2760
taggggtataa agatctcata agaaatgtaa tatgtaaaatt atatcttgct ttatgttgta 2820
aaatatacat tgtttgcgct agaatagaaa tgattttcttt tcaataaaaa gaaagaagga 2880
ctctaaaaaa aaaaaaaaaa

```

```

<210> 17
<211> 2507
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No.: 3284695CB1

```

```

<400> 17
cagagtga aa cttgtgcctg gtgaccaaag tccctccaaa gtgctcttcc ttctgggtta 60
ttcaagccaa atatctgggt ttccccctct cctcattccc tagcaaacc ccaattatctt 120
ccaagatagg agatatttcc catccccctt ctttgtaa atctcatctc ccactggaga 180
gcccaggagc ctattcctgg catggatggt ctgtccacac ttgaggctgg gcggtgtatc 240
agacccttca agcagcctgg ctggggccca ggactgagtc tggggtcagc tttcacggtc 300
gcttttccct tcctcaccac ccaccacagc ccaccttgca tgcattggca gcccctccac 360
tccagcctga gccatgtgtg cccctgcggg aggaccatt catgccagaa agctggtaac 420
tccctccag catccctgcg gaaggagtca gtttctgaga gtgtgacttt tcaaggcgaa 480
tgatggggaa ggggttccca gtccccacag tggccccacc tctgggccct gcaccagagc 540
ccttctgtgt cacggcgggc tgtgcaccca tgcacacacc tacgcacaca caaactccg 600
cactgcagta tattcttgcc aaagatttcc tttaaaagca agcactttta ctaattatta 660
ttttgtaa at gtttatcttc ttctgtcttc tccctccctg aatctatttt actgttgttt 720
attgttgaat ctgtgtgtca gccaggagag cgtgtgtctg ccttgaacat gggctgggat 780
gggaaaggg ctgggagaag atgggcaaca aagagccagg gagtcatgga catcgacagc 840
acgcagaccc cagcagggttc agtcccgtgc tggccaccagc tgtccagctg ggtgtctgga 900
gggaagaggg cagaggaggg tcatgtccct tcagctgggg gagggggcca gtgagctcca 960
cgtggctttt tcccaaaggg agcaagaggg aaggattggg cgagaaaaa atggagaggg 1020
gacctgcgaa ggaaaacagg gaggaagtga gcggtttgat cagcctgcta tcacggtgtt 1080
ctggctctct tatttagcca ggcgcttaag ggacagatac atcacatcct aagtttgga 1140
aaggcctttg acccatgtca tctgagcgtc tccctccagta gctctgaaag ctgtggacac 1200
caatggccag gattccttct cccctggttt ttgaggatcc ctgggtcttc tgagactggc 1260
caggagagg atggtggggc cagtgggtgt gtgaaagcag gaggggcagc cctcctggac 1320
aagtgtgatc cccctataaa cggctctcag gaggttagtg agtaggagat tctgccttgt 1380
tctgatgagc ctgtgcaggg gctccagggg agcatgctgt ccagggggca cagaagggtg 1440
gtgagtgtga tcaaacttag tctcactccc acttttttag tctcactcct acttttgtcc 1500
accacccctg cctcctggat cttctccac tttttttt agcttttagga cctggggaga 1560
tcctgtgagt caaggcagac acccaatcct gccccacac tcggggctct ccaagagggt 1620
ggggggcaga gtcccagagc agccctttac cccagggtcca ggccctggaa tcctgagact 1680
cgcttttctc tggccagtgg taacacagga cgtgtgtgcg catgtgcaag tgtggatgta 1740
tgtgtgtgcg tgtgttttgc tcatttcttt agggaaactg ggagtcgggg ttggaggtgc 1800
tgggcaatgg aacttcaa at tcaatgtcgc ccagcagtga ggggagtcgg gaggtgaggc 1860
ctgtaggcca accaattggg ggagtctcag cgatagccca ggtgagaagt ggttcacca 1920
gaggggcagg gtgggggcct cgggcagatc tgtccctctt ggcccctctg tcctcaa atg 1980
tccaaaatgt tggaggacct ctgttcatat cccacgcctg ggctcttgcc agcagtggag 2040
ttactgtaga gggatgtccc aagcttggtt tccaatcagt gttaaactgt ttgaaactct 2100
cctgtgtctg tgttttgttt gtgcgtgtgt gtgagagcac atcagtgtgt gcaggctgtg 2160
tttccccatt tctctctcc cttcagaccc atcattgaga acaa atgtaa gaaatccctt 2220
cccaccacc tccctgcctc ccaggccctc tgcgggggaa acaagatcac ccagcatcct 2280
tccccacccc agctgtgtat ttatatagat ggaaatatac tttatatatt gtatcatcgt 2340
gcctatagcc gctgccaccg tgtataa atc ctggtgtatg ctcttatcc tggacatgaa 2400
tgtattgtac actgacgcgt cccactcct gtacagctgc tttgtttctt tgcaatgcat 2460
tgtatggcct tataaatgat aaagttaaag aaaaaaaaaa aaaaagg 2507

```

```

<210> 18
<211> 2929
<212> DNA

```

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 3481610CB1

<400> 18

aagctcggaa	ttcggctcga	gatgggttcc	tcatcccttc	ctgctgcaaa	agaagttaac	60
aaaaaacaag	tgtgctacaa	acacaatttc	aatgcaagct	cagtttcctg	gtgttcaaaa	120
actgttgatg	tgtgttgta	ctttaccaat	gctgctaata	attcagtcctg	gagcccatct	180
atgaagctga	atctgggttcc	tggggaaaac	atcacatgcc	aggatcccgt	aataggtgtc	240
ggagagccgg	ggaaagtcac	ccagaagcta	tgccggttct	caaacgttcc	cagcagccct	300
gagagtccca	ttggcgggac	catcacttac	aaatgtgtag	gctcccagtg	ggaggagaag	360
agaaatgact	gcacatctgc	cccaataaac	agtctgctcc	agatggctaa	ggctttgatc	420
aagagccctc	ctcaggatga	gatgctccct	acatacctga	aggatctttc	tattagcata	480
ggcaaagcgg	aacatgaaat	cagctcttct	cctgggagtc	tgggagccat	tattaacatc	540
cttgatctgc	tctcaacagt	tccaacccaa	gtaaattcag	aaatgatgac	gcacgtgctc	600
tctacggtta	atatcatcct	tggcaagccc	gtcttgaaca	cctggaaggt	tttacaacag	660
caatggacca	atcagagttc	acagctacta	cattcagtg	aaagattttc	ccaagcatta	720
cagtcaggag	atagccctcc	attgtccttc	tcccaaaacta	atgtgcagat	gagcagcatg	780
gtaatcaagt	ccagccaccc	agaaacctat	caacagaggt	ttgttttccc	atactttgac	840
ctctggggca	atgtggtcat	tgacaagagc	tacctagaaa	acttgcagtc	ggattcgtct	900
attgtcacca	tggctttccc	aactctccaa	gccatccttg	ctcaggatat	ccaggaaaat	960
aactttgcag	agagcttagt	gatgacaacc	actgtcagcc	acaatacgac	tatgccattc	1020
aggatttcaa	tgacttttaa	gaacaatagc	ccttcaggcg	gcgaaacgaa	gtgtgtcttc	1080
tggaaacttca	ggcttgccaa	caacacaggg	gggtgggaca	gcagtgggtg	ctatgttgaa	1140
gaaggtgatg	gggacaatgt	cacctgtatc	tgtgaccacc	taacatcatt	ctccatcctc	1200
atgtcccctg	actccccaga	tcttagttct	ctcctgggaa	tactcctgga	tattatttct	1260
tatgttgggg	tgggcttttc	catcttgagc	ttggcagcct	gtctagtgtg	ggaagctgtg	1320
gtgtggaaat	cggtgaccaa	gaatcggact	tcttatatgc	gccacacctg	catagtgaat	1380
atcgtgcctc	cccttctggg	cgccaacacc	tgggttcattg	tggtcgctgc	catccaggac	1440
aatcgctaca	tactctgcaa	gacagcctgt	gtggctgcca	ccttcttcat	ccacttcttc	1500
tacctcagcg	tcttcttctg	gatgctgaca	ctgggcctca	tgtgtttcta	tcgcctgggt	1560
ttcattctgc	atgaaacaag	caggtccact	cagaaagcca	ttgccttctg	tcttggctat	1620
ggctgcccac	ttgccatctc	gggtcatcacg	ctgggagcca	cccagccccg	ggaagtctat	1680
acgaggaaga	atgtctgttg	gctcaactgg	gaggacacca	aggccctgct	ggctttcgcc	1740
atcccagcac	tgatcattgt	gggtgtgaac	ataaccatca	ctattgtggt	catcaccaag	1800
atcctgaggc	cttccattgg	agacaagcca	tgcaagcagg	agaagagcag	cctgtttcag	1860
atcagcaaga	gcattggggg	cctcacacca	ctcttggggc	tacttggggg	ttttggtctc	1920
accactgtgt	tcccagggac	caacctgtgt	ttccatatca	tatttgccat	cctcaatgtc	1980
ttccagggat	tattcatttt	actcttttga	tgcctctggg	atctgaaggt	acaggaagct	2040
ttgtgaata	agttttcatt	gtcgagatgg	tcttcacagc	actcaaagtc	aacatccctg	2100
ggttcatcca	cacctgtgtt	ttctatgagt	tctccaatat	caaggagatt	taacaatttg	2160
tttggtaaaa	caggaacgta	taatgtttcc	accccagaag	caaccagctc	atccctggaa	2220
aactcatcca	gtgcttcttc	gttgtctcaac	taagaacagg	ataatccaac	ctacgtgacc	2280
tcccggggac	agtggctgtg	cttttaaaaa	gagatgcttg	caaagcaatg	gggaacgtgt	2340
tctcggggca	ggtttccggg	agcagatgcc	aaaaagactt	tttcatagag	aagaggcttt	2400
cttttgtaaa	gacagaataa	aaataattgt	tatgtttctg	tttgttccct	ccccctcccc	2460
cttgtgtgat	accacatgtg	tatagtattt	aagtgaact	caagccctca	aggcccaact	2520
tctctgtcta	tattgtaata	tagaatttctg	aagagacatt	ttcacttttt	acacattggg	2580
cacaaagata	agctttgatt	aaagtagtaa	gtaaaaggct	acctaggaaa	tacttcagt	2640
aattctaaga	aggaagggaag	gaagaaagga	aggaagaaag	ggaggggaaac	agggagaaaag	2700
ggaaaaagaa	gaaaaagaga	tagatgataa	taggaacaaa	taaagacaaa	caacattaag	2760
gggcatattg	taagatttcc	atgttaatga	tctaataata	tactcagtg	ccacattttg	2820
agaatttttt	tttttaatgg	gcttcaaaaa	ttggaaaact	gtgaaagcta	agtccattgg	2880
ggggaatgga	attacttttg	ggggccagta	tctttccttt	gattgttcc		2929

<210> 19

<211> 1725

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 3722004CB1

<400> 19

```
gaggcaagaa ttcggcacga gggagagccc gcgggcggtgg gggagctcgg ggacctgcgg 60
accgggggag cccgaacgag ggggatcccc cgccggcgcc agcgaggcgg aggagcaggc 120
ggtggaggcg aggcaggaag aggagcagga cttggatggt gagaaggggc catcatcgga 180
agggcctgag gaggaggacg gagaaggctt ctccctcaaa tacagccccg ggaagctgag 240
gggaaaccag tacaagaaga tgatgaccaa agaggagctg gaggaggagc agaggattga 300
gctgacctct gacctcactt ccctgtagca agttccttag gtcctgagcc acaaatattc 360
ttgcaaatcc ttttgaactg aagaataacg aagttatcct tagcgtcctc cttaaaggctt 420
ttccttttgg catcttaaaa gcttgagaga taaaacggaa accccagaga ggagtctggg 480
caggctccca ggggtgcatg tgccctcata aatctgctga gctctagacc ctcaatcagg 540
acttgccctt tggctagcag gatcctggga acacctttgg ccctgccctg tgtagagatg 600
ttcatgtctg ttctgtggg tcaactttgtt aagctgaaga gttttaagag gtagagctca 660
gacctgggac tgggattttt cttaccactc aaacttgcta tccacacacc ctgcacacct 720
tagataaaaa gaacatttta aaagcagagt tcactttcac tccagtctcc cctcttttgc 780
cctcactgaa gccaaaccac agaagacttt gaggaatgag agacaaatga ggtagagctc 840
acctgtgctc accagctccg tcagggtggt cagccgaccc ctttccctgg gaacccact 900
tctctctgtg gctggcttgg ttgtcggggg tgagatgcc a tattgattac agggcagcaa 960
agaaccagta ccaggaattt acttgaccat tccccttatt tttcatctag aggaatctcg 1020
gattcagccc ttctattgct aagacacctt ttcactgagg ttcttaccag ctgacccaaa 1080
tctccactct gctatagcag aagcaataat gtttgcttta aaaagatttc ttgacctatg 1140
ccttttctta gaaagtttga tagattagtt agaacttcag atcatcagat cagtctcaaa 1200
tgggtttctt ggaattttat atttgacaat atttatacta taccaaaactc atttgagtt 1260
cttaggtttg ttggttaaaa cattttttta aagcagtaag tttatagaaa atgttttcat 1320
ttaatggaag gctggggaat gtccagcatc aaccctatg gcatgcattc ccagtggcct 1380
tctcatctgg gcctggaacc tttggttcag ggcttagggg agaacaggcc acatggcaac 1440
agccacacag tcattgcctt caacacagag ccacgtgtcc ccaaacagca atagtcatgc 1500
ccttgtccag gctgggatct aattgataca ataggtcgtt gactccctcc tagtagagct 1560
atctaggttt gtctggaaag tttccgaccc tggcttatag gcaccacacc tcagtactc 1620
ctcatggctt ggatctctgt attcagcctt tgttcagtcc aataaacttt gagtagatga 1680
tctcaaaaaa aaaaaaaaaa aggccggcgc aagcttattc ctttt 1725
```

<210> 20

<211> 1987

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 3948614CB1

<400> 20

```
gacggccagt gcaagctaaa attaacctc actaaaggga ataagcttgc ggccgcctgg 60
agctctcggc ctcggttcg acgacggcaa cttctcgctg ctcatccgcg cgggtggagg 120
gacggacgcg gggctgtaca cctgcaacct gcaccatcac tactgccacc tctacgagag 180
cctggccgtc cgcctggagg tcaccgacgg cccccggcc acccccgct actgggacgg 240
cgagaaggag gtgctggcgg tggcgcgcgg cgaccccgcg cttctgacct gcgtgaaccg 300
cgggcacgtg tggaccgacc ggcacgtgga ggaggtcaa caggtggtgc actgggaccg 360
gcagccgccc ggggtccgc acgaccgcgc ggaccgcctg ctggacctct acgcgtcggg 420
cgagcgccgc gcctacgggc ccctttttct gcgcgacgc gtggctgtgg gcgcggatgc 480
ctttgagcgc ggtgacttct cactgcgtat cgagccgctg gaggtcgccg acgagggcac 540
ctactcctgc cacctgcacc accattactg tggcctgcac gaacgcgcg tcttccacct 600
gacggtgcgc gaacccacg cggagccgcc cccccgggc tctccgggca acggtccag 660
ccacagcgcc gccccaggcc cagacccac actggcgcgc ggccacaacg tcatcaatgt 720
catcgtcccc gagagccgag ccacttctt ccagcagctg ggctacgtgc tggccacgct 780
gctgctcttc atcctgctac tggtcactgt cctcctggcc gccgcaggc gccgcggagg 840
ctacgaatac tcggaccaga agtcgggaaa gtcaaagggg aaggatgtta acttggcgga 900
gttcgctgtg gctgcagggg accagatgct ttacaggagt gaggacatcc agctagatta 960
caaaaacaac atcctgaagg agaggcgga gctggcccac agccccctgc ctgccaagta 1020
```

```

catcgaccta gacaaagggg tccggaagga gaactgcaaa tagggaggcc ctgggctcct 1080
ggctggggcca gcagctgcac ctctcctgtc tgtgctcctc ggggcatctc ctgatgctcc 1140
ggggctcacc ccccttccag cggctgggtc cgctttcctg gaatttgccc tgggcgtatg 1200
cagaggccgc ctccacaccc ctccccagg ggcttgggtg cagcatagcc cccacccttg 1260
cggcctttgc tcacgggttg ccctgccac ccctggcaca accaaaatcc cactgatgcc 1320
catcatgccc tcagaccctt ctgggctctg cccgctgggg gcctgaagac attcctggag 1380
gacactccca tcagaacctg gcagcccca aactggggtc agcctcaggg caggagtccc 1440
actcctccag ggctctgtct gtccggggct gggagatgtt cctggaggag gacactccca 1500
tcagaacctg gcagccttga agttggggtc agcctcggca ggagtccac tcctcctggg 1560
gtgctgcctg ccaccaagag ctccccacc tgtaccacca tgtgggactc caggcaccat 1620
ctgttctccc cagggacctg ctgacttgaa tgccagccct tgctcctctg tgttgctttg 1680
ggccacctgg ggctgcaccc cctgcccttt ctctgcccc tccctaccct agccttgctc 1740
tcagccacct tgatagtcac tgggctccct gtgacttctg accctgacac ccctcccttg 1800
gactctgcct gggctggagt ctagggtggt ggctacattt ggcttctgta ctggctgagg 1860
acaggggagg gagtgaagtt ggttgggggt ggcctgtgtt gccactctca gcaccacaca 1920
tttgcactcg ctggtggacc tgccaccatc acaataaagt ccccatctga tttttaaaaa 1980
aaaaaaa                                     1987

```

<210> 21

<211> 551

<212> PRT

<213> Homo sapiens

<220>

<221> misc feature

<223> Incyte ID No.: 627722CD1

<400> 21

```

Met Glu Glu Ala Glu Leu Val Lys Gly Arg Leu Gln Ala Ile Thr
  1          5          10          15
Asp Lys Arg Lys Ile Gln Glu Glu Ile Ser Gln Lys Arg Leu Lys
  20          25          30
Ile Glu Glu Asp Lys Leu Lys His Gln His Leu Lys Lys Lys Ala
  35          40          45
Leu Arg Glu Lys Trp Leu Leu Asp Gly Ile Ser Ser Gly Lys Glu
  50          55          60
Gln Glu Glu Met Lys Lys Gln Asn Gln Gln Asp Gln His Gln Ile
  65          70          75
Gln Val Leu Glu Gln Ser Ile Leu Arg Leu Glu Lys Glu Ile Gln
  80          85          90
Asp Leu Glu Lys Ala Glu Leu Gln Ile Ser Thr Lys Glu Glu Ala
  95          100          105
Ile Leu Lys Lys Leu Lys Ser Ile Glu Arg Thr Thr Glu Asp Ile
  110          115          120
Ile Arg Ser Val Lys Val Glu Arg Glu Glu Arg Ala Glu Glu Ser
  125          130          135
Ile Glu Asp Ile Tyr Ala Asn Ile Pro Asp Leu Pro Lys Ser Tyr
  140          145          150
Ile Pro Ser Arg Leu Arg Lys Glu Ile Asn Glu Glu Lys Glu Asp
  155          160          165
Asp Glu Gln Asn Arg Lys Ala Leu Tyr Ala Met Glu Ile Lys Val
  170          175          180
Glu Lys Asp Leu Lys Thr Gly Glu Ser Thr Val Leu Ser Ser Ile
  185          190          195
Pro Leu Pro Ser Asp Asp Phe Lys Gly Thr Gly Ile Lys Val Tyr
  200          205          210
Asp Asp Gly Gln Lys Ser Val Tyr Ala Val Ser Ser Asn His Ser
  215          220          225
Ala Ala Tyr Asn Gly Thr Asp Gly Leu Ala Pro Val Glu Val Glu
  230          235          240
Glu Leu Leu Arg Gln Ala Ser Glu Arg Asn Ser Lys Ser Pro Thr
  245          250          255

```

Glu Tyr His Glu	Pro Val Tyr Ala Asn	Pro Phe Tyr Arg Pro Thr	260	265	270
Thr Pro Gln Arg	Glu Thr Val Thr Pro	Gly Pro Asn Phe Gln Glu	275	280	285
Arg Ile Lys Ile	Lys Thr Asn Gly Leu	Gly Ile Gly Val Asn Glu	290	295	300
Ser Ile His Asn	Met Gly Asn Gly Leu	Ser Glu Glu Arg Gly Asn	305	310	315
Asn Phe Asn His	Ile Ser Pro Ile Pro	Pro Val Pro His Pro Arg	320	325	330
Ser Val Ile Gln	Gln Ala Glu Glu Lys	Leu His Thr Pro Gln Lys	335	340	345
Arg Leu Met Thr	Pro Trp Glu Glu Ser	Asn Val Met Gln Asp Lys	350	355	360
Asp Ala Pro Ser	Pro Lys Pro Arg Leu	Ser Pro Arg Glu Thr Ile	365	370	375
Phe Gly Lys Ser	Glu His Gln Asn Ser	Ser Pro Thr Cys Gln Glu	380	385	390
Asp Glu Glu Asp	Val Arg Tyr Asn Ile	Val His Ser Leu Pro Pro	395	400	405
Asp Ile Asn Asp	Thr Glu Pro Val Thr	Met Ile Phe Met Gly Tyr	410	415	420
Gln Gln Ala Glu	Asp Ser Glu Glu Asp	Lys Lys Phe Leu Thr Gly	425	430	435
Tyr Asp Gly Ile	Ile His Ala Glu Leu	Val Val Ile Asp Asp Glu	440	445	450
Glu Glu Glu Asp	Glu Gly Glu Ala Glu	Lys Pro Ser Tyr His Pro	455	460	465
Ile Ala Pro His	Ser Gln Val Tyr Gln	Pro Ala Lys Pro Thr Pro	470	475	480
Leu Pro Arg Lys	Arg Ser Glu Ala Ser	Pro His Glu Asn Thr Asn	485	490	495
His Lys Ser Pro	His Lys Asn Ser Ile	Ser Leu Lys Glu Gln Glu	500	505	510
Glu Ser Leu Gly	Ser Pro Val His His	Ser Pro Phe Asp Ala Gln	515	520	525
Thr Thr Gly Asp	Gly Thr Glu Asp Pro	Ser Leu Thr Ala Leu Arg	530	535	540
Met Arg Met Ala	Lys Leu Gly Lys Lys	Val Ile	545	550	

<210> 22

<211> 99

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 1556751CD1

<400> 22

Met Glu Ala Leu Ala	Asn Val Asn Phe	Pro Arg Lys Ser Phe Arg	1	5	10	15
Pro Glu Asp Ala Gly	Lys Glu Ser Gly	Gln Gly Gly Phe Cys	20	25	30	
Val Pro Ala Ala Arg	Pro Gln Thr Met	Val Thr Gly Pro Ser Cys	35	40	45	
Ser Ser Pro Gly Leu	Gln Asn Phe Ser	Pro Gln Arg Lys Glu Asn	50	55	60	
Arg Ala Cys Ala Cys	Trp Gln Asn Ala	Gly Pro Ala Pro Lys Asn	65	70	75	
Pro Met Cys Val Arg	Leu Lys Val Gly	Arg Pro Gln Ala Ser Gln				

Arg Lys Leu Lys Glu Thr Gly Leu Cys
 80 85 90
 95

<210> 23
 <211> 493
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No.: 2268890CD1

<400> 23
 Met Arg Pro Leu Cys Val Thr Cys Trp Trp Leu Gly Leu Leu Ala
 1 5 10 15
 Ala Met Gly Ala Val Ala Gly Gln Glu Asp Gly Phe Glu Gly Thr
 20 25 30
 Glu Glu Gly Ser Pro Arg Glu Phe Ile Tyr Leu Asn Arg Tyr Lys
 35 40 45
 Arg Ala Gly Glu Ser Gln Asp Lys Cys Thr Tyr Thr Phe Ile Val
 50 55 60
 Pro Gln Gln Arg Val Thr Gly Ala Ile Cys Val Asn Ser Lys Glu
 65 70 75
 Pro Glu Val Leu Leu Glu Asn Arg Val His Lys Gln Glu Leu Glu
 80 85 90
 Leu Leu Asn Asn Glu Leu Leu Lys Gln Lys Arg Gln Ile Glu Thr
 95 100 105
 Leu Gln Gln Leu Val Glu Val Asp Gly Gly Ile Val Ser Glu Val
 110 115 120
 Lys Leu Leu Arg Lys Glu Ser Arg Asn Met Asn Ser Arg Val Thr
 125 130 135
 Gln Leu Tyr Met Gln Leu Leu His Glu Ile Ile Arg Lys Arg Asp
 140 145 150
 Asn Ala Leu Glu Leu Ser Gln Leu Glu Asn Arg Ile Leu Asn Gln
 155 160 165
 Thr Ala Asp Met Leu Gln Leu Ala Ser Lys Tyr Lys Asp Leu Glu
 170 175 180
 His Lys Tyr Gln His Leu Ala Thr Leu Ala His Asn Gln Ser Glu
 185 190 195
 Ile Ile Ala Gln Leu Glu Glu His Cys Gln Arg Val Pro Ser Ala
 200 205 210
 Arg Pro Val Pro Gln Pro Pro Pro Ala Ala Pro Pro Arg Val Tyr
 215 220 225
 Gln Pro Pro Thr Tyr Asn Arg Ile Ile Asn Gln Ile Ser Thr Asn
 230 235 240
 Glu Ile Gln Ser Asp Gln Asn Leu Lys Val Leu Pro Pro Pro Leu
 245 250 255
 Pro Thr Met Pro Thr Leu Thr Ser Leu Pro Ser Ser Thr Asp Lys
 260 265 270
 Pro Ser Gly Pro Trp Arg Asp Cys Leu Gln Ala Leu Glu Asp Gly
 275 280 285
 His Asp Thr Ser Ser Ile Tyr Leu Val Lys Pro Glu Asn Thr Asn
 290 295 300
 Arg Leu Met Gln Val Trp Cys Asp Gln Arg His Asp Pro Gly Gly
 305 310 315
 Trp Thr Val Ile Gln Arg Arg Leu Asp Gly Ser Val Asn Phe Phe
 320 325 330
 Arg Asn Trp Glu Thr Tyr Lys Gln Gly Phe Gly Asn Ile Asp Gly
 335 340 345
 Glu Tyr Trp Leu Gly Leu Glu Asn Ile Tyr Trp Leu Thr Asn Gln
 350 355 360

Gly Asn Tyr Lys	Leu Leu Val Thr Met	Glu Asp Trp Ser Gly Arg	
	365	370	375
Lys Val Phe Ala	Glu Tyr Ala Ser Phe	Arg Leu Glu Pro Glu Ser	
	380	385	390
Glu Tyr Tyr Lys	Leu Arg Leu Gly Arg	Tyr His Gly Asn Ala Gly	
	395	400	405
Asp Ser Phe Thr	Trp His Asn Gly Lys	Gln Phe Thr Thr Leu Asp	
	410	415	420
Arg Asp His Asp	Val Tyr Thr Gly Asn	Cys Ala His Tyr Gln Lys	
	425	430	435
Gly Gly Trp Trp	Tyr Asn Ala Cys Ala	His Ser Asn Leu Asn Gly	
	440	445	450
Val Trp Tyr Arg	Gly Gly His Tyr Arg	Ser Arg Tyr Gln Asp Gly	
	455	460	465
Val Tyr Trp Ala	Glu Phe Arg Gly Gly	Ser Tyr Ser Leu Lys Lys	
	470	475	480
Val Val Met Met	Ile Arg Pro Asn Pro	Asn Thr Phe His	
	485	490	



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 9/00, C07K 14/00	A3	(11) International Publication Number: WO 00/21986 (43) International Publication Date: 20 April 2000 (20.04.00)
(21) International Application Number: PCT/US99/23315 (22) International Filing Date: 6 October 1999 (06.10.99) (30) Priority Data: 09/169,289 9 October 1998 (09.10.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/169,289 (CIP) Filed on 9 October 1998 (09.10.98) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): WALKER, Michael, G. [CA/US]; Unit 80, 1050 Borregas Avenue, Sunnyvale, CA 94089 (US). VOLKMUTH, Wayne [US/US]; 783 Roble Avenue, #1, Menlo Park, CA 94025 (US). KLINGLER, Tod, M. [US/US]; 28 Dover Court, San Carlos, CA 94070 (US). (74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 13 July 2000 (13.07.00)
(54) Title: MATRIX-REMODELING GENES		
(57) Abstract The invention provides novel matrix-remodeling genes and polypeptides encoded by those genes. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing diseases associated with matrix remodeling.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

Inter. Application No.

PCT/US 99/23315

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/00 C07K14/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MATRISIAN, L. M.: "METALLOPROTEINASES AND THEIR INHIBITORS IN MATRIX REMODELING" TRENDS IN GENETICS, vol. 6, no. 4, 1990, pages 121-125, XP000644304 * figure 2; page 123 *	1-12
A	YE, S. ET AL.: "MATRIX METALLOPROTEINASES: ..." CLINICAL SCIENCE, vol. 94, February 1998 (1998-02), pages 103-110, XP000857426 * whole disclosure *	1-12

-/--

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

27 January 2000

Date of mailing of the international search report

03.05.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

HERMANN R.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/23315

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL/GENBANK [Online] ID HSZZ74011; AC AA368885, 18 April 1997 (1997-04-18) ADAMS, M.D. ET AL.: "EST80295 Placenta I homo sapiens cDNA" XP002128855 abstract</p> <p>---</p>	2-8
X	<p>DATABASE EMBL/GENBANK [Online] ID HSW4574; AC W92457, 18 July 1996 (1996-07-18) HILLIER, L. ET AL.: "EST; Soares fetal heart NbHH19W homo sapiens cDNA clone" XP002128856 abstract</p> <p>-----</p>	2-8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/23315

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 10-12
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 10-12 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 2,7
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
See FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-3,5-7,9,10,12 (all partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 2,7

Claims 2 and 7 are -inter alia- drafted to compounds comprising as few as 18 nucleotides / 6 amino acids, which are neither defined by their exact structure, not by their exact location in the parent molecule.

Moreover, claim 2 covers

- polynucleotides of completely undefined length that hybridize to said short sequence (or the parent), and
- sequences which are complementary to the almost undefined sequences mentioned above.

Said vague structural definitions are not sufficient for a reasonable search.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-3,5-7,9,10,12 (all partially)

A polynucleotide having the SEQ ID NO 1; subject-matter of claims 2,3,5,6,9,12 relating to said polynucleotide

2. Claims: 1-12, all partially

Polypeptide with the SEQ ID NO. 21, encoding polynucleotides, and related subject-matter

3. Claims: 1-12, all partially

Polypeptide with the SEQ ID NO. 22, encoding polynucleotides, and related subject-matter

4. Claims: 1-12, all partially

Polypeptide with the SEQ ID NO. 23, encoding polynucleotides, and related subject-matter

5. Claims: 1-3,5-7,9,10,12 (all partially)

INVENTIONS 5-20:
Polynucleotides having the SEQ ID NOs. 3-5,7-10,12-20; subject-matter of claims 2,3,5,6,9,12 relating to said polynucleotide.

6. Claims: 1,3 (partially)

INVENTION 21:
Further polynucleotides and encoded polypeptides of claim 1 (PLEASE NOTE: This subject has been added for completeness only. It is considered as being at least partially anticipated (infra); Further non-unity objections may arise upon payment for subject 21)

